

## DESCRIPTION


<b>Species Reactivity</b>	Human
<b>Specificity</b>	Stains human Claudin-1 transfectants but not irrelevant transfectants.
<b>Source</b>	Monoclonal Rat IgG <sub>2A</sub> Clone # 421203
<b>Purification</b>	Protein A or G purified from hybridoma culture supernatant
<b>Immunogen</b>	CHO Chinese hamster ovary cell line transfected with human Claudin-1 Met1-Val211 Accession # O95832
<b>Conjugate</b>	Allophycocyanin Excitation Wavelength: 620-650 nm Emission Wavelength: 660-670 nm
<b>Formulation</b>	Supplied in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details.  *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

## APPLICATIONS

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Flow Cytometry	10 $\mu$ L/10 <sup>6</sup> cells	See Below

## DATA



## PREPARATION AND STORAGE

<b>Shipping</b>	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
<b>Stability &amp; Storage</b>	<b>Protect from light. Do not freeze.</b> <ul style="list-style-type: none"> <li>● 12 months from date of receipt, 2 to 8 °C as supplied.</li> </ul>

## BACKGROUND

Claudin-1 is a 23 kDa multipass membrane protein in the Claudin family of epithelial tight junction proteins. Claudin-1 is expressed by epithelial cells in a wide variety of tissues as well as in Langerhans cells and dendritic cells. It is up or downregulated in many cancers and is required for the entry of hepatitis C virus into hepatocytes. Human Claudin-1 shares 91% aa sequence identity with mouse and rat Claudin-1.