

DESCRIPTION

Species Reactivity	Mouse
Specificity	Detects mouse Jagged 2 in direct ELISAs. In direct ELISAs, no cross-reactivity with recombinant human (rh) Jagged 1, rhJagged 2, recombinant rat Jagged 1 is observed.
Source	Monoclonal Rat IgG _{2A} Clone # 746513
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Mouse myeloma cell line NS0-derived recombinant mouse Jagged 2 Met27-Leu1084 Accession # Q9QYE5
Conjugate	Alexa Fluor 405 Excitation Wavelength: 405 nm Emission Wavelength: 421 nm
Formulation	Supplied 0.2 mg/mL in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Flow Cytometry	0.25-1 µg/10 ⁶ cells	D3 mouse embryonic stem cell line

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. <ul style="list-style-type: none"> 12 months from date of receipt, 2 to 8 °C as supplied.

BACKGROUND

Mouse Jagged 2 is a 150 kDa (1) member of the Delta-Serrate-Lag-2 (DSL) family of ligands that activate LIN12/Notch proteins, and thereby regulate cell fate determinations during development (2-6). It is a type 1 transmembrane protein that is synthesized as a 1247 amino acid (aa) precursor. It contains a 23 aa signal sequence, a 1059 aa extracellular region, a 21 aa transmembrane region, and a 144 aa cytoplasmic region. The extracellular region contains four potential N-linked glycosylation sites, a DSL domain, 16 EGF-like repeats, many of which are also sites of calcium binding, and a cysteine-rich region just proximal to the transmembrane segment (3). Mouse Jagged 2 shares 93% and 90% aa sequence identity with rat and human Jagged 2, respectively. Jagged 2 is expressed highest in fetal thymus, epidermis, foregut, dorsal root ganglia, and inner ear (3). In 2-week old mice, the Jagged 2 transcript is most abundant in heart, lung, thymus, skeletal muscle, brain, and testis (3). Functionally, it is suggested that Jagged 2 engages the Notch1 pathway of signal transduction (3). It is involved in the development of the mammalian limb, branchial arches, central and peripheral nervous systems, γδ T cell lineage differentiation, and the establishment of functional natural killer cell lines (4 - 6).

References:

1. Tsai, S. *et al.* (2000) *Blood*, **96**:950.
2. Shawber, C. *et al.* (1996) *Dev. Biol.* **180**:370.
3. Luo, B. *et al.* (1997) *Mol. Cell. Biol.* **17**:6057.
4. Valsecchi, V. *et al.* (1997) *Mech. Dev.* **69**:203.
5. DeHart, S. *et al.* (2005) *Blood* **105**:3521.
6. de La Coste, A. and A.A. Freitas (2006) *Immunol. Lett.* **102**:1.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.