

DESCRIPTION

Species Reactivity	Mouse
Specificity	Detects mouse DMBT1 in direct ELISAs. In direct ELISAs, less than 5% cross-reactivity with recombinant human (rh) CD163 and rhSRCRB4D is observed.
Source	Monoclonal Rat IgG _{2B} Clone # 548031
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	<i>E. Coli</i> -derived recombinant mouse DMBT1 Trp215-Gly420 Accession # Q60997
Conjugate	Alexa Fluor 647 Excitation Wavelength: 650 nm Emission Wavelength: 668 nm
Formulation	Supplied 0.2 mg/mL in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Flow Cytometry	0.25-1 µg/10 ⁶ cells	CMT-93 mouse rectal carcinoma cell line

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. <ul style="list-style-type: none"> 12 months from date of receipt, 2 to 8 °C as supplied.

BACKGROUND

DMBT1 (deleted in malignant brain tumors 1), also called salivary agglutinin (SAG) in humans, hesperin in rabbit, ebnerin in rat, and muclin, crp-ductin or vomeroglandin in mouse, is a transmembrane or secreted ~340 kDa, 2085 amino acid (aa) protein of the scavenger receptor cysteine-rich (SRCR) family. It contains several SRCR and CUB domains and a zona pellucida motif. The region used as the immunogen (aa 215-420) contains most of SRCR 2 and 3, and shares ~74% aa identity with corresponding regions of rat or human DMBT1. Several alternately spliced, glycosylated and sulfated forms exist. DMBT1 functions in mucosal immunity by binding bacterial, viral and endogenous proteins to aid clearance by macrophages. It is also thought to play a role in epithelial cell differentiation and tumor suppression.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.