

## DESCRIPTION


|                           |  |
|---------------------------|--|
| <b>Species Reactivity</b> | Human  |
| <b>Specificity</b>        | Detects human LRRC32/GARP in ELISAs.   |
| <b>Source</b>             | Monoclonal Rat IgG <sub>2A</sub> Clone # 855151  |
| <b>Purification</b>       | Protein A or G purified from hybridoma culture supernatant   |
| <b>Immunogen</b>          | Chinese hamster ovary cell line CHO-derived recombinant human LRRC32/GARP<br>Met1-Asn627<br>Accession # Q14392   |
| <b>Conjugate</b>          | Allophycocyanin<br>Excitation Wavelength: 620-650 nm<br>Emission Wavelength: 660-670 nm  |
| <b>Formulation</b>        | Supplied in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details.<br><br>*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions. |

## APPLICATIONS

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

|                       | Recommended Concentration        | Sample    |
|-----------------------|----------------------------------|-----------|
| <b>Flow Cytometry</b> | 10 $\mu$ L/10 <sup>6</sup> cells | See Below |

## DATA



## PREPARATION AND STORAGE

|                                |  |
|--------------------------------|--|
| <b>Shipping</b>                | The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.                                  |
| <b>Stability &amp; Storage</b> | <b>Protect from light. Do not freeze.</b> <ul style="list-style-type: none"> <li>12 months from date of receipt, 2 to 8 °C as supplied.</li> </ul> |

## BACKGROUND

Leucine-rich repeat protein 32 (LRRC32), also known as GARP (glycoprotein A repetitions predominant), is an 80 kDa type I transmembrane glycoprotein (1). Mature human LRRC32 consists of a 608 amino acid (aa) extracellular domain (ECD) that contains 22 leucine-rich repeats, a 21 aa transmembrane segment, and a 14 aa cytoplasmic domain (2, 3). Within the ECD, human LRRC32 shares approximately 80% aa sequence identity with mouse and rat LRRC32. LRRC32 is widely expressed during embryogenesis and on adult platelets (4, 5). Human LRRC32 is identified as a lineage specific key receptor for human T cells. It is selectively expressed on activated FOXP3<sup>+</sup> regulatory T cells (Treg) (6-10). LRRC32 expression promotes the acquisition of a Treg phenotype including reduced cellular proliferation, reduced cytokine secretion, and the capacity to suppress the proliferation of naïve T cells (6-8). LRRC32 binds directly to the TGF-β latency associated peptide (LAP) and tethers latent TGF-β on the surface of activated Treg cells (9, 10). The presentation of TGF-β on Tregs contributes to their ability to suppress naïve T cell proliferation (11).

## References:

1. Battaglia, M. and M.G. Roncarolo (2009) *Eur. J. Immunol.* **39**:3296.
2. Ollendorff, V. *et al.* (1994) *Cell Growth Differ.* **5**:213.
3. Bella, J. *et al.* (2008) *Cell Mol Life Sci.* **65**:2307.
4. Roubin, R. *et al.* (1996) *Int. J. Dev. Biol.* **40**:545.
5. Macaulay, I.C. *et al.* (2007) *Blood* **109**:3260.
6. Wang, R. *et al.* (2008) *PLoS ONE* **3**:e2705.
7. Wang, R. *et al.* (2009) *Proc. Natl. Acad. Sci.* **106**:13439.
8. Probst-Kepper, M. *et al.* (2009) *J. Cell. Mol. Med.* **13**:3343.
9. Tran, D.Q. *et al.* (2009) *Proc. Natl. Acad. Sci.* **106**:13445.
10. Stockis, J. *et al.* (2009) *Eur. J. Immunol.* **39**:3315.
11. Vignali, D.A. *et al.* (2008) *Nat. Rev. Immunol.* **8**:523.