

#### DESCRIPTION

<b>Species Reactivity</b>	Mouse
<b>Specificity</b>	Detects mouse GM-CSF R $\alpha$ in direct ELISAs. In direct ELISAs, no cross-reactivity with recombinant human (rh) GM-CSF R alpha or rhGM-CSF R beta is observed.
<b>Source</b>	Monoclonal Rat IgG <sub>2A</sub> Clone # 698423
<b>Purification</b>	Protein A or G purified from hybridoma culture supernatant
<b>Immunogen</b>	Mouse myeloma cell line NS0-derived recombinant mouse GM-CSF R $\alpha$ Leu30-Pro327 Accession # Q00941
<b>Conjugate</b>	Alexa Fluor 647 Excitation Wavelength: 650 nm Emission Wavelength: 668 nm
<b>Formulation</b>	Supplied 0.2 mg/mL in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details.  *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

#### APPLICATIONS

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
<b>Flow Cytometry</b>	0.25-1 $\mu$ g/10 <sup>6</sup> cells	J774A.1 mouse reticulum cell sarcoma macrophage cell line

#### PREPARATION AND STORAGE

<b>Shipping</b>	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
<b>Stability &amp; Storage</b>	<b>Protect from light. Do not freeze.</b> <ul style="list-style-type: none"> <li>12 months from date of receipt, 2 to 8 °C as supplied.</li> </ul>

#### BACKGROUND

Granulocyte macrophage colony stimulating factor receptor alpha (GM-CSF R $\alpha$ ), also known as CD116, is a component of the receptor complex that mediates cellular responses to GM-CSF. GM-CSF promotes the differentiation and mobilization of granulocyte-macrophage, erythroid, megakaryocyte, and eosinophil progenitors. It enhances the activation of myeloid cell effector functions and plays a role in the development of Th1 biased immune responses, allergic inflammation, and autoimmunity (1-4). Mature mouse GM-CSF R $\alpha$  is an 80 kDa type I transmembrane glycoprotein that consists of a 298 amino acid (aa) extracellular domain (ECD) with two fibronectin type III domains and a juxtamembrane WSXWS motif, a 21 aa transmembrane segment, and a 40 aa cytoplasmic domain (5). Within the ECD, mouse GM-CSF R $\alpha$  shares approximately 33% and 58% aa sequence identity with human and rat GM-CSF R $\alpha$ , respectively. Soluble forms of the human receptor retain the ability to bind GM-CSF (6, 7). GM-CSF R $\alpha$  is expressed on hematopoietic stem cells, progenitor and differentiated cells in the myeloid lineage, vascular endothelial cells, placenta, and non-hematopoietic solid tumor cells (8). GM-CSF R $\alpha$  associates with the common beta chain/CD131 ( $\beta_c$ ), a 135 kDa transmembrane protein that is also the signal transducing component of the receptors for IL-3 and IL-5 (9, 10). Association with  $\beta_c$  converts GM-CSF R $\alpha$  from a low affinity to a high affinity receptor for GM-CSF (9-11). The shared usage of  $\beta_c$  underlies the synergism between GM-CSF, IL-3, and IL-5 in their effects on myeloid cell differentiation and activation (1, 2).

#### References:

- Martinez-Moczygemba, M. and D.P. Huston (2003) *J. Allergy Clin. Immunol.* **112**:653.
- Fleetwood, A.J. *et al.* (2005) *Crit. Rev. Immunol.* **25**:405.
- Eksioglu, E.A. *et al.* (2007) *Exp. Hematol.* **35**:1163.
- Cao, Y. (2007) *J. Clin. Invest.* **117**:2362.
- Park, L.S. *et al.* (1992) *Proc. Natl. Acad. Sci.* **89**:4295.
- Pelley, J.L. *et al.* (2007) *Exp. Hematol.* **35**:1483.
- Raines, M.A. *et al.* (1991) *Proc. Natl. Acad. Sci.* **88**:8203.
- Chiba, S. *et al.* (1990) *Cell Regul.* **1**:327.
- Kitamura, T. *et al.* (1991) *Proc. Natl. Acad. Sci.* **88**:5082.
- Hayashida, K. *et al.* (1990) *Proc. Natl. Acad. Sci.* **87**:9655.
- Hoang, T. *et al.* (1993) *J. Biol. Chem.* **268**:11881.

**PRODUCT SPECIFIC NOTICES**

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc. and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.