

## DESCRIPTION


<b>Species Reactivity</b>	Human
<b>Specificity</b>	Detects human Ly6K in direct ELISAs.
<b>Source</b>	Monoclonal Mouse IgG <sub>2A</sub> Clone # 750747
<b>Purification</b>	Protein A or G purified from hybridoma culture supernatant
<b>Immunogen</b>	Chinese hamster ovary cell line CHO-derived recombinant human Ly6K Met1-Gly138 Accession # Q17RY6
<b>Conjugate</b>	Allophycocyanin Excitation Wavelength: 620-650 nm Emission Wavelength: 660-670 nm
<b>Formulation</b>	Supplied in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details.  *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

## APPLICATIONS

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
<b>Flow Cytometry</b>	10 $\mu$ L/10 <sup>6</sup> cells	See Below

## DATA



## PREPARATION AND STORAGE

<b>Shipping</b>	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
<b>Stability &amp; Storage</b>	<b>Protect from light. Do not freeze.</b> <ul style="list-style-type: none"> <li>12 months from date of receipt, 2 to 8 °C as supplied.</li> </ul>

## BACKGROUND

Ly6K (Lymphocyte antigen 6 locus K; also CO16) is a 26-27 kDa member of the Ly-6 antigen family of molecules. It has restricted expression, being described in testis and skin. Ly6K is found in/on carcinomas, and is known to circulate in normal and tumor-patient blood. Mature human Ly6K is a 121 amino acid (aa) GPI-linked (presumed) glycoprotein. It is synthesized as a 165 aa preproprecursor that contains a 17 aa signal sequence, a 121 aa mature region (aa 18-138), and a 27 aa C-terminal propeptide. There are two potential splice variants, one that shows a 48 aa substitution, and another that shows a 33 aa substitution for aa 73-165. It is not clear if mouse Ly6K is a true molecular ortholog of human Ly6K. In any event, over aa 18-138, human Ly6K shares 39% aa identity with mouse Ly6K.