

DESCRIPTION


Species Reactivity	Mouse/Rat
Specificity	Detects mouse and rat DARC in Western blots.
Source	Polyclonal Sheep IgG
Purification	Antigen Affinity-purified
Immunogen	E. coli-derived recombinant mouse DARC Met1-Pro61, Ala115-Cys127, Ser186-Lys205, Tyr264-Asn285 Accession # NP_034175
Conjugate	Phycoerythrin Excitation Wavelength: 488 nm Emission Wavelength: 565-605 nm
Formulation	Supplied in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Flow Cytometry	10 μ L/10 ⁶ cells	See Below

DATA



PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. <ul style="list-style-type: none"> 12 months from date of receipt, 2 to 8 °C as supplied.

BACKGROUND

DARC (Duffy Antigen Receptor for Chemokines; also CD234) is a 40-46 kDa glycoprotein member of the Duffy family of silent heptahelical chemokine receptors. It is expressed in liver and on select neurons, erythrocytes and the endothelium of postcapillary venules. Unlike traditional chemokine receptors, DARC cannot signal through G-proteins as it lacks a DRYLAIVHA cytoplasmic motif. DARC has three potential functions: first, it binds circulating inflammatory-type chemokines, serving as a repository for future chemokine release; second, it acts as a vehicle by which chemokines are transported from the abluminal to the luminal side of endothelium; and third, it complexes with signal-transducing chemokine receptors to create a nonsignaling heterodimer. Mouse DARC is 334 amino acids (aa) in length. It contains a 62 aa N-terminal extracellular region, and a 28 aa C-terminal cytoplasmic tail. There is one potential splice variant that shows a 42 aa substitution for aa 133-334. Collectively, over the four extracellular domains (aa 1-62, 115-127, 186-205, 264-285), mouse DARC shares 52% and 75% aa identity with human and rat DARC, respectively.