

DESCRIPTION

Species Reactivity	Mouse
Specificity	Stains mouse IL-7 R α /CD127 transfectants but not irrelevant transfectants in flow cytometry.
Source	Monoclonal Rabbit IgG Clone # 1140A
Purification	Protein A or G purified from cell culture supernatant
Immunogen	NSO mouse myeloma cell line transfected with mouse IL-7 R α /CD127 Glu21-Asp239 Accession # P16872
Conjugate	Alexa Fluor 700 Excitation Wavelength: 675-700 nm Emission Wavelength: 723 nm
Formulation	Supplied 0.2 mg/mL in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Flow Cytometry	0.25-1 μ g/10 ⁶ cells	Mouse splenocytes

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage **Protect from light. Do not freeze.**

- 12 months from date of receipt, 2 to 8 °C as supplied.

BACKGROUND

Interleukin 7 Receptor alpha (IL-7 R α), also known as CD127, is a 75 kDa hematopoietin receptor superfamily member that plays an important role in lymphocyte differentiation, proliferation, and survival (1, 2). Mature mouse IL-7 R α consists of a 219 amino acid (aa) extracellular domain (ECD) with one fibronectin type III domain and a WSxWS motif, a 25 aa transmembrane segment, and a 195 aa cytoplasmic domain (3). Within the ECD, mouse IL-7 R α shares 67% and 79% aa sequence identity with human and rat IL-7 R α , respectively. IL-7 R α associates with the common γ chain (γ_c) to form the functional high affinity IL-7 receptor complex (4). The γ_c is also a subunit of the receptors for IL-2, -4, -9, -15, and -21. Human and mouse IL-7 show cross-species activity through the IL-7 receptor (3, 5). IL-7 R α is expressed on double negative (CD4⁻CD8⁻) and CD4⁺ or CD8⁺ single positive T cells as well as on CD8⁺ memory T cells and their precursors (6, 7). It is expressed early in B cell development, prior to the appearance of surface IgM (6). In mouse, IL-7 activation of IL-7 R α is critical for both T cell and B cell lineage development (8). In human it is required for T cell but not for B cell development (9). IL-7 induces the down regulation and shedding of cell surface IL-7 R α (10). IL-7 R α additionally associates with TSLP R to form the functional receptor for thymic stromal lymphopoietin (11, 12). TSLP indirectly regulates T cell development by modulating dendritic cell activation (2, 13). Knockout of TSLP R in mice provokes minor changes in B and T cell development compared to those seen with IL-7 R α deletion (8, 14). The complexity of IL-7 R α biology is suggested by the competition between IL-7 and TSLP for receptor binding and by the ability of IL-7 R α to form functional complexes with SCF R and HGF R (11, 12, 15, 16).

References:

1. Mazzucchelli, R. and S.K. Durum, 2007, Nat. Rev. Immunol. 7:144.
2. Liu, Y.-J. *et al.* (2007) Annu. Rev. Immunol. 25:193.
3. Goodwin, R.G. *et al.* (1990) Cell 60:941.
4. Noguchi, M. *et al.* (1993) Science 262:1877.
5. Barata, J.T. *et al.* (2006), Exp. Hematol. 34:1133.
6. Sudo, T. *et al.* (1993) Proc. Natl. Acad. Sci. 90:9125.
7. Kaech, S.M. *et al.* (2003) Nat. Immunol. 4:1191.
8. Peschon, J.J. *et al.* (1994) J. Exp. Med. 180:1955.
9. Prieyl, J.A. and T.W. LeBien (1996) Proc. Natl. Acad. Sci. 93:10348.
10. Vranjkovic, A. *et al.* (2007) Int. Immunol. 19:1329.
11. Park, L.S. *et al.* (2000) J. Exp. Med. 192:659.
12. Pandey, A. *et al.* (2000) Nat. Immunol. 1:59.
13. Reche, P.A. *et al.* (2001) J. Immunol. 167:336.
14. Al-Shami, A. *et al.* (2004) J. Exp. Med. 200:159.
15. Jahn, T. *et al.* (2007) Blood 110:1840.
16. Lai, L. *et al.* (2006) Blood 107:1776.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.