

# Mouse Common $\gamma$ Chain/IL-2 R $\gamma$ Alexa Fluor® 647-conjugated Antibody

Monoclonal Rat IgG<sub>2B</sub> Clone # 644346

Catalog Number: FAB7842R

100  $\mu$ g

## DESCRIPTION

<b>Species Reactivity</b>	Mouse
<b>Specificity</b>	Detects mouse Common $\gamma$ Chain/IL-2 R $\gamma$ in direct ELISAs. In direct ELISAs, no cross-reactivity with recombinant mouse (rm) IL-2 R $\alpha$ , rmlL-2 R $\beta$ , recombinant human (rh) IL-2 R $\gamma$ , rhIL-15 R $\alpha$ , recombinant rat IL-2 R $\alpha$ , or recombinant canine IL-2 R $\alpha$ is observed.
<b>Source</b>	Monoclonal Rat IgG <sub>2B</sub> Clone # 644346
<b>Purification</b>	Protein A or G purified from hybridoma culture supernatant
<b>Immunogen</b>	<i>S. frugiperda</i> insect ovarian cell line Sf21-derived recombinant mouse Common $\gamma$ Chain/IL-2 R $\gamma$ Trp23-Ala263 (predicted) Accession # P34902
<b>Conjugate</b>	Alexa Fluor 647 Excitation Wavelength: 650 nm Emission Wavelength: 668 nm
<b>Formulation</b>	Supplied 0.2 mg/mL in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details.  *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

## APPLICATIONS

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.

	Recommended Concentration	Sample
<b>Flow Cytometry</b>	0.25-1 $\mu$ g/10 <sup>6</sup> cells	CD3+ mouse splenocytes

## PREPARATION AND STORAGE

<b>Shipping</b>	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
<b>Stability &amp; Storage</b>	<b>Protect from light. Do not freeze.</b> <ul style="list-style-type: none"> <li>12 months from date of receipt, 2 to 8 °C as supplied.</li> </ul>

## BACKGROUND

The Common gamma Chain, also known as CD132, is a transmembrane protein belonging to the hematopoietin receptor family.  $\gamma$ c is a signaling component of the functional receptor complexes for IL-2, IL-4, IL-7, IL-9, and IL-15.

## PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.