

DESCRIPTION

| | |
|---------------------------|--|
| Species Reactivity | Rat |
| Specificity | Detects rat CXCR3 in ELISAs. |
| Source | Recombinant Monoclonal Mouse IgG _{2B} Clone # 868013R |
| Purification | Protein A or G purified from cell culture supernatant |
| Immunogen | NS0 mouse myeloma cell line transfected with rat CXCR3 Accession # Q9J119 |
| Conjugate | Alexa Fluor 488 Excitation Wavelength: 488 nm Emission Wavelength: 515-545 nm |
| Formulation | Supplied 0.2 mg/mL in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions. |

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

| | Recommended Concentration | Sample |
|-----------------------|----------------------------------|---|
| Flow Cytometry | 0.25-1 µg/10 ⁶ cells | NS0 Cell Line Transfected with Rat CXCR3 and eGFP |

PREPARATION AND STORAGE

| | |
|--------------------------------|--|
| Shipping | The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below. |
| Stability & Storage | Protect from light. Do not freeze. <ul style="list-style-type: none"> 12 months from date of receipt, 2 to 8 °C as supplied. |

BACKGROUND

CXCR3, also known as CD183, is an approximately 41 kDa (predicted) 7-transmembrane spanning receptor for the chemokines CXCL9, CXCL10, and CXCL11. It is expressed on activated T cells, B cells, and NK cells during inflammation. It is additionally upregulated on solid tumor cells, tumor endothelium, and cancer stem cells. CXCR3 plays a role in leukocyte recruitment to sites of inflammation. It also contributes to Th1 biased responses during autoimmune diseases rheumatoid arthritis, systemic erythematosis, and type 1 diabetes. Rat CXCR3 shares 86% and 96% amino acid sequence identity with human and mouse CXCR3, respectively.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.