

DESCRIPTION

Species Reactivity	Mouse
Specificity	Detects mouse DC-SIGN/CD209 in direct ELISAs.
Source	Monoclonal Rat IgG _{2B} Clone # 902404
Purification	Protein A or G purified from cell culture supernatant
Immunogen	NS0 mouse myeloma cell line transfected with mouse DC-SIGN/CD209 Met1-Lys238 Accession # Q91ZX1
Conjugate	Alexa Fluor 647 Excitation Wavelength: 650 nm Emission Wavelength: 668 nm
Formulation	Supplied 0.2 mg/mL in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Flow Cytometry	0.25-1 µg/10 ⁶ cells	CHO Chinese hamster ovary cell line transfected with mouse DC-SIGN/CD209

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. <ul style="list-style-type: none"> ● 12 months from date of receipt, 2 to 8 °C as supplied.

BACKGROUND

Mouse Dendritic Cell-specific ICAM-3 Grabbing Non-integrin (DC-SIGN)/CD209, also known as CD209 Antigen-like Protein A, is a member of the C-type lectin family (1). Mouse DC-SIGN/CD209 is a 33 kDa, 238 amino acid (aa) type II transmembrane protein (2). The extracellular region contains a Ca²⁺-dependent carbohydrate-binding lectin domain (2). In addition to the full-length mouse DC-SIGN/CD209, a second, truncated splice variant has been reported. DC-SIGN/CD209 is not well conserved between mouse and human with the extracellular domain sharing only 63% aa identity. The DC-SIGN/CD209 lectin domain binds mannose oligosaccharides on pathogens including HIV as well as self glycoproteins including ICAMs (2-4). DC-SIGN/CD209 is expressed on dendritic cells (DC) and inflammatory macrophages and contributes to antigen presentation (2, 5, 6).

References:

1. Liu, W. *et al.* (2004) *J. Biol. Chem.* **279**:18748.
2. Caminschi, I. *et al.* (2001) *Mol. Immunol.* **38**:365.
3. Curtis, B.M. *et al.* (1992) *Proc. Natl. Acad. Sci. USA* **89**:8356.
4. Anthony, R.M. *et al.* (2008) *Proc. Natl. Acad. Sci. USA* **105**:19571.
5. Geijtenbeek, T.B. *et al.* (2000) *Cell* **100**:575.
6. Garcia-Vallejo, J.J. and Y. van Kooyk (2013) *Trends Immunol.* **34**:482.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.