

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human ICAM-4 in direct ELISAs.
Source	Monoclonal Mouse IgG _{2B} Clone # 729632
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Chinese hamster ovary cell line CHO-derived recombinant human ICAM-4 Ala23-Ala240 Accession # Q14773
Conjugate	Alexa Fluor 647 Excitation Wavelength: 650 nm Emission Wavelength: 668 nm
Formulation	Supplied 0.2 mg/mL in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Flow Cytometry	0.25-1 µg/10 ⁶ cells	Human red blood cells

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. <ul style="list-style-type: none"> ● 12 months from date of receipt, 2 to 8 °C as supplied.

BACKGROUND

ICAM-4 (Intercellular Adhesion Molecule 4), also known as Landsteiner-Wiener Glycoprotein and CD242, is a 42 kDa member of the ICAM family, Ig superfamily of proteins. It is expressed on erythrocytes and erythroblasts, and serves as a receptor for LFA-1, Mac-1, and CD11c/CD18, plus α4β1 and alpha-V containing integrins. ICAM-4 is suggested to bind to Mac-1 on macrophages, allowing for its phagocytosis in senescence. Mature human ICAM-4 is a 249 amino acid (aa) type I transmembrane glycoprotein. It possesses a 218 aa extracellular region (aa 23-240) that contains two C2-type Ig-like domains (aa 62-124 and 146-217), and a 10 aa C-terminal cytoplasmic tail. ICAM-4 may form 85 kDa homodimers. There are three potential isoform variants. One shows a five aa substitution for aa 233-271, a second contains a 15 aa substitution for aa 14-29, and a third possesses a 141 aa substitution for aa 132-271. Over aa 31-240, human ICAM-4 shares 71% aa identity with mouse ICAM-4.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.