

DESCRIPTION


Species Reactivity	Human
Specificity	Stains human EMR3 transfected but not irrelevant transfected cells in flow cytometry.
Source	Monoclonal Mouse IgG _{2B} Clone # 908235
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	NS0 mouse myeloma cell line transfected with human EMR3 Met1-Tyr652 Accession # NP_115960
Conjugate	Allophycocyanin Excitation Wavelength: 620-650 nm Emission Wavelength: 660-670 nm
Formulation	Supplied in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Flow Cytometry	10 μ L/10 ⁶ cells	See Below

DATA



PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. <ul style="list-style-type: none"> 12 months from date of receipt, 2 to 8 °C as supplied.

BACKGROUND

EMR3 is a member of the class B seven-span transmembrane (TM7) receptor family. It is expressed predominantly by cells of the immune system, with highest levels in neutrophils, monocytes and macrophages. It may play a role in myeloid-myeloid interactions during immune and inflammatory responses. EMR3 upregulation in glioblastoma multiforme (GBM) is associated with poor survival and may be a potential therapeutic target.