

#### DESCRIPTION

<b>Species Reactivity</b>	Mouse
<b>Specificity</b>	Detects mouse CXCR1 transfectants but not CXCR2 transfectants by flow cytometry.
<b>Source</b>	Monoclonal Rabbit IgG Clone # 1122A
<b>Purification</b>	Protein A or G purified from cell culture supernatant
<b>Immunogen</b>	Mouse CXCR1/IL-8 RA Peptide Accession # Q810W6
<b>Conjugate</b>	Alexa Fluor 405 Excitation Wavelength: 405 nm Emission Wavelength: 421 nm
<b>Formulation</b>	Supplied 0.2 mg/mL in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details.  *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

#### APPLICATIONS

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	<b>Recommended Concentration</b>	<b>Sample</b>
<b>Flow Cytometry</b>	0.25-1 µg/10 <sup>6</sup> cells	Human whole blood monocytes

#### PREPARATION AND STORAGE

<b>Shipping</b>	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
<b>Stability &amp; Storage</b>	<b>Protect from light. Do not freeze.</b> <ul style="list-style-type: none"> <li>12 months from date of receipt, 2 to 8 °C as supplied.</li> </ul>

#### BACKGROUND

CXCR1, also known as IL-8 RA and CD181, is an approximately 60 kDa 7TM glycoprotein that functions as a receptor for the chemokine CXCL8/IL-8. It is expressed on neutrophils, monocytes, CD8 T cells, FoxP3+ CD4 Treg cells, mast cells, neuronal and glial cells, vascular endothelial cells, and melanoma. CXCR1 forms homodimers and heterodimers with CXCR2/IL-8 RB. It can be cleaved from neutrophils in the lungs of cystic fibrosis patients to release fragments that promote CXCL8 production from airway epithelial cells. CXCR1 mediates neutrophil activation and chemotaxis to sites of inflammation as well as angiogenesis and melanoma invasiveness. Mouse CXCR1 shares 65% and 89% amino acid sequence identity with human and rat CXCR1, respectively.

#### PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.