

MATERIAL DATA SHEET

Nedd8 E1 Activating Enzyme Inhibitor (NAE inhibitor), Synthetic Cat. # I-502

NAE inhibitor contains the active component of MLN4924, a cell-permeable, small molecule inhibitor of Nedd8 activating enzyme (NAE). By inhibiting the Neddylation pathway this molecule disrupts cullin-RING ligase activity, leading to inhibition of proliferation or apoptotic death of various human tumor cells through the deregulation of S-phase DNA synthesis. Mutations in UBA3, a gene encoding one of the subunits of Nedd8 activating enzyme, have been reported to cause resistance to MLN4924 in some tumor cell lines.

Product Information

Quantity: 1 mg

FW: 444

Structure:

Physical/Chemical Characteristics

Solubility: Provided as a 20 mM stock solution in DMSO

Purity: > 99% by HPLC

Activity: Depending on cell line, typical IC50 values in 72 hour assay are $\leq 1 \mu M$.

Use & Storage

Storage: Store lyophilized material at -20°C. Store reconstituted material at -20°C for

up to 3 months. Avoid multiple freeze/thaw cycles.

Literature

References: Lin, J.J., et al. (2010) Cancer Res. **70**: 10310-10320

Soucy, T.A., et al. (2009) Nature 458: 732-737

For Laboratory Research Use Only, Not For Use in Humans

Rev: 09/25/12

840 Memorial Drive, Cambridge, MA 02139 Phone: 617-576-2210 FAX: 617-492-3565 <u>www.bostonbiochem.com</u>