

DESCRIPTION

Species Reactivity	Mouse
Specificity	Detects mouse IL-17F in direct ELISAs and Western blots. In direct ELISAs and Western blots, no cross-reactivity with recombinant mouse (rm) IL-17, rmlL-17B, C, D, E, or recombinant human IL-17F is observed.
Source	Monoclonal Rat IgG _{2A} Clone # 316016
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	<i>E. coli</i> -derived recombinant mouse IL-17F Arg29-Ala161 Accession # Q7TNI7
Conjugate	Alexa Fluor 700 Excitation Wavelength: 675-700 nm Emission Wavelength: 723 nm
Formulation	Supplied 0.2 mg/mL in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Intracellular Staining by Flow Cytometry	0.25-1 µg/10 ⁶ cells	Mouse splenocytes treated with PMA, Ca ²⁺ ionomycin, Recombinant Human TGF-β (Catalog # 240-B), and Recombinant Mouse IL-6 (Catalog # 406-ML) fixed with paraformaldehyde and permeabilized with methanol and saponin

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. <ul style="list-style-type: none"> 12 months from date of receipt, 2 to 8 °C as supplied.

BACKGROUND

The Interleukin 17 (IL-17) family proteins, comprised of six members (IL-17, IL-17B through IL-17F), are secreted, structurally related proteins that share a conserved cysteine-knot fold near the C-terminus, but have considerable sequence divergence at the N-terminus. With the exception of IL-17B, which exists as a non-covalently linked dimer, all IL-17 family members are disulfide-linked dimers. IL-17 family proteins are pro-inflammatory cytokines that induce local cytokine production and are involved in the regulation of immune functions (1, 2).

Mouse IL-17F cDNA encodes a 153 amino acid (aa) protein with a putative 20 aa signal peptide. Among IL-17 family members, IL-17F is most closely related to IL-17 sharing approximately 46% aa sequence identity. Mouse and human IL-17F share 55% sequence identity. IL-17F is expressed in activated CD4⁺ T cells and activated monocytes. Two receptors (IL-17 R, and IL-17B R), which are activated by IL-17 family members have been identified. In addition, at least three additional type I transmembrane receptors with homology to IL-17 R, including IL-17 RL (IL-17 RC), IL-17 RD, and IL-17 RE, have also been reported (1, 2, 5). The functions for IL-17 RC, D, and E are not known. Purified IL-17 R and IL-17B R do not bind IL-17F with high-affinity *in vitro*. However, binding of IL-17F is detected in cells transfected with IL-17 R, raising the possibility that a co-receptor may be required for IL-17F signaling through IL-17 R (4). The biological activities mediated by IL-17F are similar to those of IL-17. IL-17F stimulates production of IL-6, IL-8, G-CSF, and regulates cartilage matrix turnover by increasing matrix release and inhibiting new matrix synthesis (4). IL-17F also inhibits angiogenesis and induces production of IL-2, TGF-β, and monocyte chemoattractant protein-1 in endothelial cells (3).

References:

1. Aggarwal, S. and A.L. Gurney (2002) *J. Leukoc. Biol.* **71**:1.
2. Moseley, T.A. *et al.* (2003) *Cytokine & Growth Factor Rev.* **14**:155.
3. Starnes, T. *et al.* (2001) *J. Immunol.* **167**:4137.
4. Hurst, S.D. *et al.* (2002) *J. Immunol.* **169**:443.
5. Haudenschild, D. *et al.* (2002) *J. Biol. Chem.* **277**:4309.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.