

DESCRIPTION


Species Reactivity	Human
Specificity	Detects TGF-β1 from human, mouse, rat, and other species in direct ELISAs and Western blots. In sandwich ELISAs, less than 2% cross-reactivity with recombinant human (rh) TGF-β3 and recombinant amphibian TGF-β5 and no cross-reactivity with recombinant porcine TGF-β2 or rhTGF-β2 is observed.
Source	Monoclonal Mouse IgG ₁ Clone # 9016
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Chinese hamster ovary cell line CHO-derived recombinant TGF-β1 and latent TGF-β1
Conjugate	Allophycocyanin Excitation Wavelength: 620-650 nm Emission Wavelength: 660-670 nm
Formulation	Supplied in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the [Technical Information](#) section on our website.

	Recommended Concentration	Sample
Intracellular Staining by Flow Cytometry	10 μL/10 ⁶ cells	See Below

DATA



PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. <ul style="list-style-type: none"> ● 12 months from date of receipt, 2 to 8 °C as supplied.

BACKGROUND

TGF-β1, -2, and -3 are a closely related group of proteins (70-80% sequence homology) that are produced by many cell types and function as growth and differentiation factors. The active forms of TGF-β1, -2, and -3 are disulfide-linked homodimers.