

# Human VEGF Alexa Fluor® 488-conjugated Antibody

Monoclonal Mouse IgG<sub>2A</sub> Clone # 23410

Catalog Number: IC2931G 100 TESTS


DESCRIPTION			
Species Reactivity	Human		
Specificity	Detects human VEGF in flow cytometry.		
Source	Monoclonal Mouse IgG <sub>2A</sub> Clone # 23410		
Purification	Protein A or G purified from ascites		
Immunogen	E. coli-derived recombinant human VEGF		
Conjugate	Alexa Fluor 488 Excitation Wavelength: 488 nm Emission Wavelength: 515-545 nm		
Formulation	Supplied in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details.		
	*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet		

#### **APPLICATIONS**

Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.

	Recommended Concentration	Sample
Intracellular Staining by Flow Cytometry	5 μL/10 <sup>6</sup> cells	See Below

#### DATA



Detection of VEGF in U937 Human Cell Line by Flow Cytometry. Activated U937 human histiocytic lymphoma cell line was stained with Mouse Anti-Human VEGF Alexa Fluor® 488-conjugated Monoclonal Antibody (Catalog # IC2931G, filled histogram) or isotype control antibody (Catalog # IC003G, open histogram). To facilitate intracellular staining, cells were fixed with Flow Cytometry Fixation Buffer (Catalog # FC004) and permeabilized with Flow Cytometry Permeabilization/Wash Buffer I (Catalog # FC005). View our protocol for Staining Intracellular Molecules.

(SDS) for additional information and handling instructions.

### PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage

Protect from light. Do not freeze.

12 months from date of receipt, 2 to 8 °C as supplied

## BACKGROUND

Vascular Endothelial Growth Factor (VEGF) is a soluble protein secreted by a wide variety of cell types. It binds to the receptor tyrosine kinases VEGF R1 (Flt-1) and VEGF R2 (Flk-1). VEGF stimulates vascular endothelial cell proliferation and is a potent inducer of angiogenesis. Several VEGF isoforms occur resulting from alternative mRNA splicing.

## PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.

