

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human IL-17/IL-17A in direct ELISAs. In direct ELISAs, approximately 12% cross-reactivity with recombinant canine IL-17/IL-17A is observed and 25%-50% reactivity with recombinant human (rh) IL-17A/IL-17F heterodimer is observed. No cross-reactivity with recombinant mouse IL-17/IL-17A, rhIL-17B, rhIL-17C, rhIL-17D, rhIL-17E, or rhIL-17F is observed.
Source	Monoclonal Mouse IgG _{2B} Clone # 41809
Purification	Protein A or G purified from ascites
Immunogen	<i>E. coli</i> -derived recombinant human IL-17/IL-17A Ile20-Ala155 Accession # Q16552
Conjugate	Allophycocyanin Excitation Wavelength: 620-650 nm Emission Wavelength: 660-670 nm
Formulation	Supplied in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.


APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Intracellular Staining by Flow Cytometry	10 μ L/10 ⁶ cells	See Below

DATA

Intracellular Staining by Flow Cytometry



Detection of IL-17/IL-17A in Human PBMCs by Flow Cytometry. Human peripheral blood mononuclear cells (PBMCs) either (A) untreated or (B) treated with 50 ng/mL PMA, 200 ng/mL Calcium Ionomycin, 200 ng/mL LPS, 20 ng/mL Recombinant Human IL-23 (Catalog # [1290-IL](#)), and 40 ng/mL Recombinant Human IL-6 (Catalog # [206-IL](#)) overnight to induce Th17 development were stained with Mouse Anti-Human IL-17/IL-17A APC-conjugated Monoclonal Antibody (Catalog # IC317A) and Mouse Anti-Human CD3 ϵ PE-conjugated Monoclonal Antibody (Catalog # [FAB100P](#)). Quadrant markers were set based on control antibody staining (Catalog # [IC0041A](#)). To facilitate intracellular staining, cells were fixed with Flow Cytometry Fixation Buffer (Catalog # [FC004](#)) and permeabilized with Flow Cytometry Permeabilization/Wash Buffer I (Catalog # [FC005](#)). View our protocol for [Staining Intracellular Molecules](#).

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. <ul style="list-style-type: none"> 12 months from date of receipt, 2 to 8 °C as supplied.

BACKGROUND

IL-17/IL-17A is a pro-inflammatory cytokine secreted by activated T cells. It is the prototype member of the IL-17 family that also includes IL-17B, C, D, E, and F.