

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human FoxP1 in direct ELISAs and Western blots.
Source	Monoclonal Mouse IgG ₁ Clone # 837016
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	<i>E. coli</i> -derived recombinant human FoxP1 Lys548-Glu677 Accession # Q9H334
Conjugate	Alexa Fluor 647 Excitation Wavelength: 650 nm Emission Wavelength: 668 nm
Formulation	Supplied 0.2 mg/mL in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Intracellular Staining by Flow Cytometry	0.25-1 µg/10 ⁶ cells	MCF-7 human breast cancer cell line fixed with 4% paraformaldehyde and permeabilized with methanol

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. <ul style="list-style-type: none"> 12 months from date of receipt, 2 to 8 °C as supplied.

BACKGROUND

Forkhead Box P1 (FOXP1) is a member of the FOX family of transcription factors. FoxP1 has been implicated in cardiac, lung, and lymphocyte development. FoxP1 knock out mice die at embryonic day 14.5 due to heart valve and outflow tract abnormalities. FoxP1 contains both a DNA binding domain as well as protein-protein interaction domains. FoxP1 can homo or heterodimerize with FoxP2 and FoxP4, with dimerization necessary for DNA binding. FoxP1 shows both oncogenic and tumor suppressive characteristics. Overexpression in lymphomas leads to poor prognosis, but loss of FoxP1 in breast cancer also implicates a poor prognosis. Human isoforms of 489 to 677 amino acids contain alternate sequences within the first 60 amino acids and/or deletion of amino acids.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.