

DESCRIPTION

Species Reactivity	Mouse
Specificity	Detects mouse IL-21 in Western blots. Does not cross-react with recombinant human IL-21, recombinant mouse (rm) IL-15, or rmlIL-12.
Source	Monoclonal Rat IgG _{2B} Clone # 149204
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	<i>E. coli</i> -derived recombinant mouse IL-21 Pro25-Ser146 Accession # Q9ES17
Conjugate	Alexa Fluor 594 Excitation Wavelength: 590 nm Emission Wavelength: 617 nm
Formulation	Supplied 0.2 mg/mL in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Intracellular Staining by Flow Cytometry	0.25-1 µg/10 ⁶ cells	Mouse splenocytes treated with PMA and Ca ²⁺ ionomycin, fixed with paraformaldehyde, and permeabilized with saponin

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. <ul style="list-style-type: none"> 12 months from date of receipt, 2 to 8 °C as supplied.

BACKGROUND

Interleukin-21 (IL-21) and its receptor appear to play important roles in the regulation of the immune system. IL-21 is most related to IL-2, IL-4, and IL-15. IL-21 R, also called NILR (novel interleukin receptor), is a type I cytokine receptor with 4 conserved cysteine residues and an extracellular WSXWS motif. It is most closely related to IL-2 Rβ and IL-4 Rα. Mouse IL-21 is a 146 amino acid (aa) residue protein with a 24 aa signal peptide. Mouse and human IL-21 share 57% aa identity. IL-21 is expressed by activated T cells. Although not fully elucidated, the IL-2 Rγ (γ_c) chain appears to play a role in IL-21 R signaling. The IL-21/IL-21 R interaction appear to play important roles in B and T cell proliferation after antigen stimulation and NK cell maturation.

References:

1. Parrish-Novak, *et al.* (2000) *Nature* **408**:57.
2. Ozaki, K. *et al.* (2000) *Proc. Natl. Acad. Sci. USA* **97**:11439.
3. Dumoutier, L. *et al.* (2000) *Proc. Natl. Acad. Sci. USA* **97**:10144.
4. Asao, H. *et al.* (2001) *J. Immunol.* **167**:1.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.