

DESCRIPTION


Species Reactivity	Human/Mouse
Specificity	Detects human and mouse ROR γ t/RORC2 in direct ELISAs. In direct ELISAs, no cross-reactivity with human and mouse RORC.
Source	Monoclonal Mouse IgG _{2B} Clone # 600380
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Human ROR γ t peptide Met1-Arg10 Accession # P51449
Conjugate	PerCP (Peridinin-chlorophyll Protein Complex) Excitation Wavelength: 482 and 564 nm Emission Wavelength: 675 nm
Formulation	Supplied in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Intracellular Staining by Flow Cytometry	10 μ L/10 ⁶ cells	See Below

DATA



PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. <ul style="list-style-type: none"> 12 months from date of receipt, 2 to 8 °C as supplied.

BACKGROUND

Retinoic acid-related Orphan Receptor gamma (ROR γ , TOR, RORC; NR1F3) is a member of the orphan nuclear receptor family. ROR γ is expressed in the muscle, thymus, testis, pancreas, prostate, heart, and liver. ROR γ plays a role in thymocyte development and homeostasis. RORs bind to DNA as monomers on half-site elements with 5' A/T-rich extensions. An N-terminal isoform of ROR γ , ROR γ t, has been shown to be specifically expressed in the thymus.