

DESCRIPTION

Specificity	Selected for use as mouse IgG ₁ isotype control. In direct ELISAs, does not cross-react with recombinant human (rh) 6Ckine, recombinant mouse (rm) 6Ckine, rhACT-II, rhBLC/BCA-1, rmBLC, rmC10, recombinant rat (rr) CINC-1, rrCINC-2α, rrCINC-2β, rrCINC-3, rhCKβ8-1, recombinant viral (rv) CMV UL146, rmCRG-2, rhENA-78, rhEotaxin, rmEotaxin, rhEotaxin-2, rmEotaxin-2, rhEotaxin-3, rhFractalkine, rmFractalkine, rrFractalkine, rhGCP-2, rmGCP-2, rhGROα, rhGROβ, rhGROγ, rhHCC-1, rhI-309, rhI-Linck, rhIL-8, recombinant porcine IL-8, rhIP-10, rhI-TAC, rmJE, rmKC, rhLeukotactin-1, rhLymphotactin, rmLymphotactin, rmMARC, rhMCP-1, rhMCP-2, rhMCP-3, rhMCP-4, rmMCP-5, rvMCV type 2, rhMDC, rmMDC, rhMIG, rmMIG, rhMIP-1α, rmMIP-1α, rhMIP-1β, rhMIP-1δ, rhMIP-1γ, rmMIP-2, rhMIP-3α, rmMIP-3α, rrMIP-3α, rhMIP-3β, rmMIP-3β, rvMIP-1, rvMIP-II, rvMIP-III, rhMPIF-1, rhNAP-2, rhPARC, rhRANTES, rmRANTES, rhSDF-1α, rmSDF-1α, rhSDF-1β, rhTARC, rmTARC, rmTCA-3, rhTeck, or rmTeck.
Source	Monoclonal Mouse IgG ₁ κ Clone # 11711R
Purification	Protein A or G purified from cell culture supernatant
Immunogen	KLH
Formulation	Supplied as a solution in PBS. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 μm filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.

Negative control for use in conjunction with R&D Systems antibodies. Use at the same concentration as the detection antibody.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C, as supplied. ● 1 month, 2 to 8 °C under sterile conditions after opening. ● 6 months, -20 to -70 °C under sterile conditions after opening.