

Mouse IL-18 Rα/IL-1 R5 Antibody

Monoclonal Rat IgG₁ Clone # 112614

Catalog Number: MAB1216


DESCRIPTION			
Species Reactivity	Mouse		
Specificity	Detects mouse IL-18 Ra/IL-1 R5 in direct ELISAs and Western blots. In direct ELISAs and Western blots, no cross-reactivity with recombinant mouse (rm) IL-1 RI, rmIL-1 RII, recombinant human (rh) IL-1 R3, rhIL-1 R4, recombinant rat (rr) IL-1 R6, rmIL-1 R7, rhIL-1 R rhIL-1 R9, or rmSIGIRR is observed.		
Source	Monoclonal Rat IgG ₁ Clone # 112614		
Purification	Protein A or G purified from hybridoma culture supernatant		
Immunogen	Mouse myeloma cell line NS0-derived recombinant mouse IL-18 Rα/IL-1 R5 Ser19-Gly326 Accession # Q61098		
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS.		

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.

	Recommended Concentration	Sample
Western Blot	1 μg/mL	Recombinant Mouse IL-18 Rα/IL-1 R5
Flow Cytometry	2.5 μg/10 ⁶ cells	See Below

DATA



Detection of IL-18 Rα/IL-1 R5 in Mouse Splenocytes by Flow Cytometry. Mouse splenocytes were stained with Rat Anti-Mouse IL-18 Rα/IL-1 R5 Monoclonal Antibody (Catalog # MAB1216, filled histogram) or isotype control antibody (Catalog # MAB005, open histogram), followed by Phycoerythrin-conjugated Anti-Rat IgG F(ab')₂ Secondary Antibody (Catalog # F0105B).

PREPARATION AND STORAGE

Reconstitution Reconstitute at 0.5 mg/mL in sterile PBS

Shipping The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.

*Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C

- 12 months from date of receipt, -20 to -70 °C as supplied.
- 1 month, 2 to 8 °C under sterile conditions after reconstitution.
- 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

IL-18 R, also known as IL-18 R alpha, is a member of the IL-1 receptor family and has been designated IL-1 R5

