

#### DESCRIPTION


<b>Species Reactivity</b>	Human
<b>Specificity</b>	Detects human IL-17F in direct ELISAs.
<b>Source</b>	Recombinant Monoclonal Mouse IgG <sub>2B</sub> Clone # 775608R
<b>Purification</b>	Protein A or G purified from cell culture supernatant
<b>Immunogen</b>	Mouse myeloma cell line NS0-derived recombinant human IL-17F Gly21-Thr153 Accession # Q96PD4
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 µm filtered solution in PBS.

#### APPLICATIONS

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the Technical Information section on our website.

<b>ELISA</b>	This antibody functions as an ELISA capture antibody when paired with Mouse Anti-Human IL-17F Monoclonal Antibody (Catalog # <a href="#">MAB13353</a> ).  <i>This product is intended for assay development on various assay platforms requiring antibody pairs. We recommend the Human IL-17F DuoSet ELISA Kit (Catalog # <a href="#">DY1335B-05</a>) for convenient development of a sandwich ELISA.</i>
--------------	--

#### DATA



#### PREPARATION AND STORAGE

<b>Reconstitution</b>	Reconstitute at 0.5 mg/mL in sterile PBS.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
<b>Stability &amp; Storage</b>	<b>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</b> <ul style="list-style-type: none"> <li>• 12 months from date of receipt, -20 to -70 °C as supplied.</li> <li>• 1 month, 2 to 8 °C under sterile conditions after reconstitution.</li> <li>• 6 months, -20 to -70 °C under sterile conditions after reconstitution.</li> </ul>

**BACKGROUND**

The Interleukin 17 (IL-17) family proteins, comprising six members (IL-17A through IL-17F), are secreted, structurally related proteins that share a conserved cystine-knot fold near the C-terminus, but have considerable sequence divergence at the N-terminus. With the exception of IL-17B, which exists as a non-covalently linked dimer, all IL-17 family members are disulfide-linked dimers. IL-17 family proteins are pro-inflammatory cytokines that induce local cytokine production and are involved in the regulation of immune functions (1, 2).

Human IL-17F cDNA encodes a 163 amino acid (a) protein with a putative 30 aa signal peptide. Among IL-17 family members, IL-17F is most closely related to IL-17A (approximately 44% aa sequence homology), but shares only limited sequence homology (16-30%) with IL-17B, C, D and E. Human and mouse IL-17F share 55% sequence identity. IL-17F is expressed in activated CD4<sup>+</sup> T-cells and activated monocytes. Five receptors (IL-17 RA, B, C, D and E) have been identified (5). Although the ligands for IL-17 RD and E are not known yet, it is reported that IL-17 RA binds IL-17A, and IL-17 RB binds IL-17B and IL-17E. IL-17 RC binds IL-17A and IL-17F with similarly high affinity and functions as a receptor for both IL-17A and IL-17F (5, 6). The biological activities mediated by IL-17F are similar to those of IL-17. IL-17F stimulates production of IL-6, IL-8, G-CSF, and regulates cartilage matrix turnover by increasing matrix release and inhibiting new matrix synthesis (4). IL-17F also inhibits angiogenesis and induces production of IL-2, TGF-β, and monocyte chemoattractant protein-1 in endothelial cells (3).

**References:**

1. Aggarwal, S. and A.L. Gurney (2002) J. Leukoc. Biol. **71**:1.
2. Moseley, T.A. *et al.* (2003) Cytokine & Growth Factor Rev. **14**:155.
3. Starnes, T. *et al.* (2001) J. Immunol. **167**:4137.
4. Shen, F. & S. L. Gaffen (2008) Cytokine **41**:92.
5. Kuestner, R.E. *et al.* (2007) J. Immunol. **179**:5462.