

DESCRIPTION

Species Reactivity	Human/Mouse/Bovine
Specificity	Detects bovine, human, and mouse insulin.
Source	Monoclonal Rat IgG _{2A} Clone # 182410
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	<i>E. coli</i> -derived recombinant human Insulin
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS.


APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Immunocytochemistry	8-25 µg/mL	See Below
Intracellular Staining by Flow Cytometry	2.5 µg/10 ⁶ cells	βTC-6 mouse beta cell insulinoma cell line fixed with paraformaldehyde and permeabilized with saponin

DATA

Immunocytochemistry



Insulin in βTC-6 Mouse Cell Line. Insulin was detected in immersion fixed βTC-6 mouse beta cell insulinoma cell line using Human/Mouse/Bovine Insulin Monoclonal Antibody (Catalog # MAB1417) at 10 µg/mL for 3 hours at room temperature. Cells were stained using the NorthernLights™ 557-conjugated Anti-Rat IgG Secondary Antibody (red; Catalog # NL013) and counterstained with DAPI (blue). View our protocol for [Fluorescent ICC Staining of Cells on Coverslips](#).

PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.5 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

Insulin is a disulfide-linked heterodimeric protein secreted by the pancreatic Islets of Langerhans. Mature insulin is generated by the proteolytic removal of a peptide from proinsulin. It is involved in the regulation of glucose metabolism through interactions with the Insulin Receptor.