

DESCRIPTION

| | |
|---------------------------|--|
| Species Reactivity | Mouse |
| Specificity | Detects mouse Fetuin A/AHSG in ELISAs. In sandwich immunoassays, no cross-reactivity or interference was observed with recombinant human Fetuin A, rhTGF-β1, recombinant mouse (rm) BMP-4, rmMMP-2, recombinant rat MMP-2, or rmMMP-9 is observed. |
| Source | Monoclonal Rat IgG _{2B} Clone # 207440 |
| Purification | Protein A or G purified from hybridoma culture supernatant |
| Immunogen | Mouse myeloma cell line NS0-derived recombinant mouse Fetuin A/AHSG Ala19-Ile345 Accession # P29699 |
| Formulation | Lyophilized from a 0.2 μm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 μm filtered solution in PBS. |

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

| Mouse Fetuin A/AHSG Sandwich Immunoassay | | Reagent |
|---|---------------|---|
| ELISA Capture | 2-8 μg/mL | Mouse Fetuin A/AHSG Antibody (Catalog # MAB15632) |
| ELISA Detection | 0.5-2.0 μg/mL | Mouse Fetuin A/AHSG Biotinylated Antibody (Catalog # BAM15631) |
| Standard | | Recombinant Mouse Fetuin A/AHSG (Catalog # 1563-PI) |

PREPARATION AND STORAGE

| | |
|--------------------------------|--|
| Reconstitution | Reconstitute at 0.5 mg/mL in sterile PBS. |
| Shipping | The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C |
| Stability & Storage | Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution. |

BACKGROUND

Mouse Fetuin A, also known as α₂-Heremans-Schmid glycoprotein, is encoded by the AHSG gene. It has been also called "countertrypsin" because of its ability to inhibit trypsin (1). It is a major plasma protein and a member of the cystatin superfamily of protease inhibitors (2, 3). It is expressed by hepatocytes, the principal cell source, and by monocyte/macrophages (4). The major form of plasma Fetuin A corresponds to two disulfide bond-linked chains derived from the single chain (5). Fetuin-A has a number of functions. It is a negative acute-phase protein with normal circulating levels in adults (300-600 μg/mL), which fall significantly (30-50%) during injury and infection (5). It enhances entry of cationic inhibitors into macrophages (6). It inhibits both insulin receptor autophosphorylation and undesirable calcification (7, 8).

References:

1. Yamamoto, K. and H. Sinohara (1993) J. Biol. Chem. **268**:17750.
2. Kelleman, J. *et al.* (1989) J. Biol. Chem. **264**:14121.
3. Dziegielewska, K.M. *et al.* (1990) J. Biol. Chem. **265**:4354.
4. Dziegielewska, K.M. *et al.* (1996) Histochem. Cell Biol. **106**:319.
5. Gejyo, F. and K. Schmid (1981) Biochim. Biophys. Acta **671**:78.
6. Wang, H. *et al.* (1998) Proc. Natl. Acad. Sci. USA **95**:14429.
7. Mathews, S.T. *et al.* (2000) Mol. Cell Endocrinol. **164**:87.
8. Schäfer, C. *et al.* (2003) J. Clin. Invest. **112**:357.