

## DESCRIPTION

<b>Species Reactivity</b>	Human/Mouse
<b>Specificity</b>	Detects human and mouse SOX2 in Western blots.
<b>Source</b>	Monoclonal Mouse IgG <sub>2A</sub> Clone # 245610
<b>Purification</b>	Protein A or G purified from hybridoma culture supernatant
<b>Immunogen</b>	<i>E. coli</i> -derived recombinant human SOX2 Gly135-Met317 Accession # P48431
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS.


## APPLICATIONS

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
<b>Western Blot</b>	0.2 µg/mL	See Below
<b>Immunocytochemistry</b>	8-25 µg/mL	See Below
<b>Intracellular Staining by Flow Cytometry</b>	2.5 µg/10 <sup>6</sup> cells	NTera-2 human embryonic carcinoma cell line fixed with paraformaldehyde and permeabilized with saponin
<b>Simple Western</b>	4 µg/mL	See Below


## DATA

### Immunocytochemistry




**E-Cadherin and SOX2 in BG01V Human Stem Cells.** E-Cadherin and SOX2 were detected in BG01V human embryonic stem cells using 10 µg/mL Goat Anti-Human E-Cadherin Antigen Affinity-purified Polyclonal Antibody (Catalog # AF648) and 10 µg/mL Mouse Anti-Human/Mouse SOX2 Monoclonal Antibody (Catalog # MAB2018). Cells were incubated with primary antibodies for 3 hours at room temperature. Cells were stained for E-Cadherin using the NorthernLights™ 557-conjugated Anti-Goat IgG Secondary Antibody (green; Catalog # NL001) and for SOX2 using the NorthernLights 493-conjugated Anti-Mouse Secondary Antibody (red; Catalog # NL009). Cells were counterstained with DAPI (blue). View our protocol for [Fluorescent ICC Staining of Cells on Coverslips](#).

### Immunocytochemistry




**SOX2 in BG01V Human Embryonic Stem Cells.** SOX2 was detected in immersion fixed BG01V human embryonic stem cells using Mouse Anti-Mouse SOX2 Monoclonal Antibody (Catalog # MAB2018) at 10 µg/mL for 3 hours at room temperature. Cells were stained using the NorthernLights™ 557-conjugated Anti-Mouse IgG Secondary Antibody (yellow; Catalog # NL007) and counterstained with DAPI (blue). View our protocol for [Fluorescent ICC Staining of Cells on Coverslips](#).

### Western Blot



**Detection of Human SOX2 by Western Blot.** Western blot shows lysates of BG01V human embryonic stem cells and NTera-2 human testicular embryonic carcinoma cell line. PVDF membrane was probed with 0.2 µg/mL of Mouse Anti-Human/Mouse SOX2 Monoclonal Antibody (Catalog # MAB2018) followed by HRP-conjugated Anti-Mouse IgG Secondary Antibody (Catalog # HAF007). This experiment was conducted under reducing conditions and using [Immunoblot Buffer Group 1](#).

### Simple Western



**Detection of Human SOX2 by Simple Western™.** Simple Western lane view shows lysates of BG01V human embryonic stem cells, loaded at 0.2 mg/mL. A specific band was detected for SOX2 at approximately 50 kDa (as indicated) using 4 µg/mL of Mouse Anti-Human/Mouse SOX2 Monoclonal Antibody (Catalog # MAB2018). This experiment was conducted under reducing conditions and using the 12-230 kDa separation system.



## PREPARATION AND STORAGE

**Reconstitution** Reconstitute at 0.5 mg/mL in sterile PBS.

**Shipping** The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.  
\*Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C

**Stability & Storage** Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

- 12 months from date of receipt, -20 to -70 °C as supplied.
- 1 month, 2 to 8 °C under sterile conditions after reconstitution.
- 6 months, -20 to -70 °C under sterile conditions after reconstitution.

## BACKGROUND

SOX2 belongs to the SOX (SRY-like HMG box) family of transcription factors with diverse roles in development. SOX2 functions in specifying the first three lineages present at implantation and in regulating proliferation and differentiation in the developing peripheral nervous system (1-6).

### References:

1. Graham, V. *et al.* (2003) *Neuron* **39**:749.
2. Avilion, A.A. *et al.* (2003) *Genes Dev.* **17**:126.
3. Kishi, M. *et al.* (2000) *Development* **127**:791.
4. Yuan, H. *et al.* (1995) *Genes Dev.* **9**:2635.
5. Uwanogho, D. *et al.* (1995) *Mech. Dev.* **49**:23.
6. Stevanovic, M. (2003) *Mol. Biol. Rep.* **30**:127.