

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human M-CSF R in ELISAs. In ELISAs, this antibody does not cross-react with recombinant human (rh) M-CSF, rhGM-CSF, rhPDGF R α , rhPDGF R β , or rmM-CSF.
Source	Monoclonal Mouse IgG ₁ Clone # 61715
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Mouse myeloma cell line NS0-derived recombinant human M-CSF R Ile20-Glu512 (Pro54Ala) Accession # P07333.2
Formulation	Lyophilized from a 0.2 μ m filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 μ m filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

Human M-CSF R Sandwich Immunoassay		Reagent
ELISA Capture	2-8 μ g/mL	Human M-CSF R Antibody (Catalog # MAB3292)
ELISA Detection	0.1-0.4 μ g/mL	Human M-CSF R Biotinylated Antibody (Catalog # BAF329)
Standard		Recombinant Human M-CSF R Fc Chimera (Catalog # 329-MR)

PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.5 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> • 12 months from date of receipt, -20 to -70 °C as supplied. • 1 month, 2 to 8 °C under sterile conditions after reconstitution. • 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

M-CSF receptor, the product of the *c-fms* proto-oncogene, is a member of the type III subfamily of receptor tyrosine kinases that also includes receptors for SCF and PDGF. These receptors each contain five immunoglobulin-like domains in their extracellular domain (ECD) and a split kinase domain in their intracellular region (1-4). M-CSF receptor is expressed primarily on cells of the monocyte/macrophage lineage, dendritic cells, stem cells and in the developing placenta (1). Human M-CSF receptor cDNA encodes a 972 amino acid (aa) type I membrane protein with a 19 aa signal peptide, a 493 aa extracellular region containing the ligand-binding domain, a 25 aa transmembrane domain and a 435 aa cytoplasmic domain. The human M-CSF R ECD shares 60%, 64%, 72%, 75%, 75%, and 76% aa identity with mouse, rat, bovine, canine, feline and equine M-CSF R, respectively. Activators of protein kinase C induce TACE/ADAM17 cleavage of the M-CSF receptor, releasing the functional ligand-binding extracellular domain (5). M-CSF binding induces receptor homodimerization, resulting in transphosphorylation of specific cytoplasmic tyrosine residues and signal transduction (6). The intracellular domain of activated M-CSF R binds more than 150 proteins that affect cell proliferation, survival, differentiation and cytoskeletal reorganization. Among these, PI3Kinase, P42/44 ERK and c-Cbl are key transducers of M-CSF R signals (3, 4). M-CSF R engagement is continuously required for macrophage survival and regulates lineage decisions and maturation of monocytes, macrophages, osteoclasts and DC (3, 4). M-CSF R and integrin $\alpha\beta_3$ share signaling pathways during osteoclastogenesis and deletion of either causes osteopetrosis (7, 8). In the brain, microglia expressing increased M-CSF R are concentrated with Alzheimers $\alpha\beta$ peptide, but their role in pathogenesis is unclear (9, 10).

References:

1. deParseval, N. *et al.* (1993) *Nucleic Acids Res.* **21**:750.
2. Rothwell, V.M. and L.R. Rohrschneider (1987) *Oncogene Res.* **1**:311.
3. Chitu, V. and E.R. Stanley (2006) *Curr. Opin. Immunol.* **18**:39.
4. Ross, F.P. and S.L. Teitelbaum (2005) *Immunol. Rev.* **208**:88.
5. Rovida, E. *et al.* (2001) *J. Immunol.* **166**:1583.
6. Yeung, Y. *et al.* (1998) *J. Biol. Chem.* **273**:17128.
7. Dai, X. *et al.* (2002) *Blood* **99**:111.
8. Faccio, R. *et al.* (2003) *J. Clin. Invest.* **111**:749.
9. Li, M. *et al.* (2004) *J. Neurochem.* **91**:623.
10. Mitrasinovic, O.M. *et al.* (2005) *J. Neurosci.* **25**:4442.