

## DESCRIPTION


<b>Species Reactivity</b>	Human
<b>Specificity</b>	Detects human Catalase in ELISAs. In direct ELISAs, no cross-reactivity with recombinant mouse Catalase or recombinant human Serpin C1 is observed.
<b>Source</b>	Monoclonal Mouse IgG <sub>1</sub> Clone # 724810
<b>Purification</b>	Protein A or G purified from hybridoma culture supernatant
<b>Immunogen</b>	<i>E. coli</i> -derived recombinant human Catalase Met1-Leu527 Accession # P04040
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS.

## APPLICATIONS

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the [Technical Information](#) section on our website.

	Recommended Concentration	Sample
<b>Western Blot</b>	0.5 µg/mL	See Below

## DATA



## PREPARATION AND STORAGE

<b>Reconstitution</b>	Reconstitute at 0.5 mg/mL in sterile PBS.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
<b>Stability &amp; Storage</b>	<b>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</b> <ul style="list-style-type: none"> <li>● 12 months from date of receipt, -20 to -70 °C as supplied.</li> <li>● 1 month, 2 to 8 °C under sterile conditions after reconstitution.</li> <li>● 6 months, -20 to -70 °C under sterile conditions after reconstitution.</li> </ul>

## BACKGROUND

Cells have evolved complex mechanisms to maintain redox balance and defend against oxidative stress. Catalase is a tetrameric enzyme comprised of four 60 kDa subunits. Catalase is typically localized in the peroxisome where it functions as an antioxidant, protecting cells from damage due to oxidative stress. Catalase converts reactive oxygen species, such as H<sub>2</sub>O<sub>2</sub>, into water and O<sub>2</sub>. Human Catalase shares 89% homology to mouse and rat Catalase. The cells redox environment can serve as an important signaling switch or trigger to initiate a number of cellular processes, including gene expression, differentiation, proliferation and apoptosis.