

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human ErbB3/Her3 in direct ELISAs.
Source	Monoclonal Mouse IgG _{2B} Clone # 526806
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	<i>E. coli</i> -derived recombinant human ErbB3/Her3 Asn953-Gly1090 Accession # P21860
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS.


APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Immunohistochemistry	8-25 µg/mL	See Below

DATA

Immunohistochemistry



ErbB3/Her3 in Human Lung. ErbB3/Her3 was detected in immersion fixed paraffin-embedded sections of human lung array using Mouse Anti-Human ErbB3/Her3 Monoclonal Antibody (Catalog # MAB3483) at 15 µg/mL overnight at 4 °C. Tissue was stained using the Anti-Mouse HRP-DAB Cell & Tissue Staining Kit (brown; Catalog # CTS002) and counterstained with hematoxylin (blue). Lower panel shows a lack of labeling if primary antibodies are omitted and tissue is stained only with secondary antibody followed by incubation with detection reagents. View our protocol for [Chromogenic IHC Staining of Paraffin-embedded Tissue Sections](#).

PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.5 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

ErbB3, also called Her3 (human epidermal growth factor receptor 3), is a type I membrane glycoprotein that is a member of the ErbB family of tyrosine kinase receptors. ErbB family members serve as receptors for the epidermal growth factor (EGF) family of growth factors. Among ErbB family members, ErbB3 is unique in that it contains a defective kinase domain. ErbB3 is expressed in keratinocytes, melanocytes, skeletal muscle cells, embryonic myoblasts and Schwann cells. Monomeric ErbB3 serves as a low affinity receptor for the heregulins (HRG). ErbB3 heterodimerizes with ErbB2 to form a high affinity receptor complex. In contrast, ErbB3 homodimerization or heterodimerization with ErbB4 forms a low affinity heregulin-binding complex. Because ErbB3 contains a defective kinase domain, the kinase domain of ErbB2 is responsible for initiating the tyrosine phosphorylation signal through the heterodimeric receptor. It has been found that a discrete three amino acid signal in the ErbB3 cytoplasmic domain is critical for transactivation of ErbB2. The cytoplasmic domain of ErbB3 also contains six consensus binding motifs for the SH2 domain of the regulatory p85 subunit of phosphoinositide 3-kinase (PI 3-kinase, PI3K) as well as one proline-rich consensus binding motif for the SH3 domain of p85. Human ErbB3 consists of 1342 amino acids (aa) with a 19 aa signal sequence, a 624 aa extracellular domain, a 21 aa transmembrane region, and a 678 aa cytoplasmic domain. ErbB3 appears to play roles in development, cancer, communication at the neuromuscular junction and regulation of cell growth and differentiation.

References:

1. Kraus, M.H. *et al.* (1989) Proc. Natl. Acad. Sci. **86**:9193.
2. Plowman, G.D. *et al.* (1990) Proc. Natl. Acad. Sci. USA **87**:4905.
3. Carraway, K.L. 3rd *et al.* (1994) J. Biol. Chem. **269**:14303.
4. Emkey, R. and C.R. Kahn (1997) J. Biol. Chem. **272**:31172.
5. Sundaresan, S. *et al.* (1998) Endocrinology **139**:4756.
6. Hellyer, N.J. *et al.* (1998) Biochem. J. **333**:757.
7. Schaefer, G. *et al.* (1999) J. Biol. Chem. **274**:859.
8. Hellyer, N.J. *et al.* (2001) J. Biol. Chem. **276**:42153.
9. Schlessinger, J. (2000) Cell **103**:211.
10. Daly, R.J. (1999) Growth Factors **16**:255.