

DESCRIPTION


Species Reactivity	Human
Specificity	Detects human CCL17/TARC in ELISAs. Does not cross-react with recombinant human (rh) CCL2, 3, 4, 5, 7, 8, 11, 13, 16, 18, 19, 20, 21, 22, 23, 24, 25, rmCCL2, 3, 4, 5, 7, 9/10/MIP-1 γ, 11, 12, 21, 22, 25, or rCCL20.
Source	Monoclonal Mouse IgG ₁ Clone # 54026
Purification	Protein A or G purified from ascites
Immunogen	<i>E. coli</i> -derived recombinant human CCL17/TARC Ala24-Ser94 Accession # Q92583.1
Endotoxin Level	<0.10 EU per 1 μg of the antibody by the LAL method.
Formulation	Lyophilized from a 0.2 μm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 μm filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

Human CCL17/TARC Sandwich Immunoassay	Reagent
ELISA Capture	2-8 μg/mL Human CCL17/TARC Antibody (Catalog # MAB364)
ELISA Detection	0.1-0.4 μg/mL Human CCL17/TARC Biotinylated Antibody (Catalog # BAF364)
Standard	Recombinant Human CCL17/TARC (Catalog # 364-DN)
Neutralization	Measured by its ability to neutralize CCL17/TARC-induced chemotaxis in the BaF3 mouse pro-B cell line transfected with human CCR4. The Neutralization Dose (ND ₅₀) is typically 0.4-2.0 μg/mL in the presence of 10 ng/mL Recombinant Human CCL17/TARC.

DATA



PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.5 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

CCL17 is a novel CC chemokine identified using a signal sequence trap method. CCL17 cDNA encodes a highly basic 94 amino acid (aa)residue precursor protein with a 23 aa residue signal peptide that is cleaved to generate the 71 aa residue mature secreted protein. Among CC chemokine family members, CCL17 has approximately 24-29% amino acid sequence identity with RANTES, MIP-1α, MIP-1β, MCP-1, MCP-2, MCP-3, and I-309. The gene for human CCL17 has been mapped to chromosome 16q13 rather than chromosome 17 where the genes for many human CC chemokines are clustered. CCL17 is constitutively expressed in thymus, and at a lower level in lung, colon, and small intestine. CCL17 is also transiently expressed in stimulated peripheral blood mononuclear cells. Recombinant CCL17 has been shown to be chemotactic for T cell lines but not monocytes or neutrophils. CCL17 was identified to be a specific functional ligand for CCR4, a receptor that is selectively expressed on T cells.

References:

1. Imai, T. *et al.* (1997) *J. Biol. Chem.* **272**:15036.
2. Imai, T. *et al.* (1996) *J. Biol. Chem.* **271**:21514.
3. Nomiyama, H. *et al.* (1997) *Genomics* **40**:211.

