

DESCRIPTION


Species Reactivity	Human
Specificity	Detects human DC-LAMP in direct ELISAs. In direct ELISAs, no cross-reactivity with recombinant human LAMP is observed.
Source	Monoclonal Mouse IgG _{2A} Clone # 530217
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Mouse myeloma cell line NS0-derived recombinant human DC-LAMP Asp21-Thr381 Accession # EAW78338
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Western Blot	2 µg/mL	See Below

DATA



PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.5 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

DC-LAMP (dendritic cell lysosome-associated membrane protein; also CD208 and LAMP3) is a member of the LAMP family of proteins. Mature human DC-LAMP is a 389 amino acid (aa) type I transmembrane protein. It can be variably glycosylated, and apparent sizes from 54kDa to 90 kDa have been reported. It has a 254 aa luminal N-terminus and a short 14 aa cytoplasmic tail. The molecule is found in type II pneumocytes, interdigitating DC, and various tumors. It may play a role in normal lysosome and endosome function. Over aa 21-381, human DC-LAMP shares 72% and 55% aa sequence identity with dog and mouse DC-LAMP, respectively.