

## DESCRIPTION


<b>Species Reactivity</b>	Human
<b>Specificity</b>	Detects human Enolase 2/Neuron-specific Enolase in direct ELISAs and Western blots.
<b>Source</b>	Monoclonal Mouse IgG <sub>2B</sub> Clone # 522601
<b>Purification</b>	Protein A or G purified from hybridoma culture supernatant
<b>Immunogen</b>	<i>E. coli</i> -derived recombinant human Enolase 2/Neuron-specific Enolase Met1-Leu434 Accession # P09104
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution in PBS and NaCl with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS.

## APPLICATIONS

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
<b>Western Blot</b>	2 µg/mL	See Below
<b>Immunohistochemistry</b>	8-25 µg/mL	Immersion fixed paraffin-embedded sections of human brain (cortex)
<b>Immunoprecipitation</b>	25 µg/mL	Cell lysates spiked with Recombinant Human Enolase 2, <a href="#">see our available Western blot detection antibodies</a>

## DATA



## PREPARATION AND STORAGE

<b>Reconstitution</b>	Reconstitute at 0.5 mg/mL in sterile PBS.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
<b>Stability &amp; Storage</b>	<b>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</b> <ul style="list-style-type: none"> <li>● 12 months from date of receipt, -20 to -70 °C as supplied.</li> <li>● 1 month, 2 to 8 °C under sterile conditions after reconstitution.</li> <li>● 6 months, -20 to -70 °C under sterile conditions after reconstitution.</li> </ul>

## BACKGROUND

Enolase 2 (2-phospho-D-glycerate hydrolyase; also Neuron-specific Enolase, NSE, neural enolase and  $\gamma$ -enolase) is a 46 kDa member of the Enolase family of enzymes. It is expressed in developing neurons and glia, is known to catalyze the generation of phosphoenolpyruvate, and is suggested to possess neurotrophic activity for neurons, likely through an extracellular mechanism. Human Enolase 2 is 434 amino acids (aa) in length. The enzymatic site spans most of the length of the molecule. Enolase 2 exists as both a noncovalently-linked homodimer, or heterodimer with  $\alpha$ -enolase. Full-length human Enolase 2 shares 99% aa identity with both mouse and canine Enolase 2. It shares 83% aa identity with human enolases 1 and 3.