

DESCRIPTION




Species Reactivity	Human
Specificity	Detects human ZNF281 in direct ELISAs and Western blots.
Source	Monoclonal Mouse IgG ₁ Clone # 801706
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	<i>E. coli</i> -derived recombinant human ZNF281 Asn681-Arg895 Accession # Q9Y2X9
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Western Blot	1 µg/mL	See Below
Immunocytochemistry	8-25 µg/mL	See Below
Simple Western	50 µg/mL	See Below

DATA



PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.5 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> • 12 months from date of receipt, -20 to -70 °C as supplied. • 1 month, 2 to 8 °C under sterile conditions after reconstitution. • 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

ZNF281 (Zinc Finger protein 28), also known as ZBP99 and GZP1, is a 100 kDa member of the Krueppel C2H2-type Zn-finger family of proteins. It is found in embryonic stem cells, placenta, and lymphocytes, and appears to be necessary for the maintenance of pluripotency in stem cells. ZNF281 both represses (STAT3) and activates (Nanog) multiple genes, and is known to form a complex with Nanog, Oct4 and Sox2. Human ZNF281 is 895 amino acids in length. It contains one Gly-rich region (aa 4-37), a poly-Pro motif (aa 90-96) and four consecutive C2H2-type zinc finger domains (aa 261-367). There are phosphorylation sites at Ser255, 395, 484, 785 and 807. Potential alternate start sites exist at Met296 and Met434. Over aa 681-895, human ZNF281 shares 99% aa identity with mouse ZNF281.