

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human Gastrokine 1 in direct ELISAs. In direct ELISAs, no cross-reactivity with recombinant mouse Gastrokine 1 is observed.
Source	Monoclonal Mouse IgG ₁ Clone # 665701
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	<i>E. coli</i> -derived recombinant human Gastrokine 1 Asn35-Asn199 Accession # Q9NS71
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS.


APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Immunohistochemistry	8-25 µg/mL	See Below

DATA

Immunohistochemistry



Gastrokine 1 in Human Stomach.

Gastrokine 1 was detected in immersion fixed paraffin-embedded sections of human stomach using Mouse Anti-Human Gastrokine 1 Monoclonal Antibody (Catalog # MAB6395) at 15 µg/mL overnight at 4 °C. Before incubation with the primary antibody, tissue was subjected to heat-induced epitope retrieval using Antigen Retrieval Reagent-Basic (Catalog # CTS013). Tissue was stained using the Anti-Mouse HRP-DAB Cell & Tissue Staining Kit (brown; Catalog # CTS002) and counterstained with hematoxylin (blue). Specific staining was localized to gastric glands. View our protocol for [Chromogenic IHC Staining of Paraffin-embedded Tissue Sections](#).

PREPARATION AND STORAGE

Reconstitution	Sterile PBS to a final concentration of 0.5 mg/mL.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

Gastrokine 1 (GKN1; also CA11 and AMP-18) is an 18 kDa member of the CA11 protein family. It has limited expression, being restricted to mucous secreting pyloric atrial epithelium. Gastrokine 1 appears to promote epithelial proliferation and migration, and induce the formation of tight junctions between epithelial cells. Mature human Gastrokine 1 is 165 amino acids (aa) in length. Based on the SwissProt sequence, it possesses one BRICHOS domain (aa 54-150) that contains a mitogenic sequence (aa 97-117). There is one potential alternative start site 14 aa upstream of the standard site that does not appear to be utilized. Over aa 21-185 (aa 35-199 of SwissProt), human Gastrokine 1 shares 65% aa identity with mouse Gastrokine 1.