

## DESCRIPTION


<b>Species Reactivity</b>	Mouse
<b>Specificity</b>	Detects mouse MINA in ELISAs. In direct ELISAs, approximately 50% cross-reactivity with recombinant human MINA is observed.
<b>Source</b>	Monoclonal Rat IgG <sub>2A</sub> Clone # 768826
<b>Purification</b>	Protein A or G purified from hybridoma culture supernatant
<b>Immunogen</b>	<i>E. coli</i> -derived recombinant mouse MINA Met1-Gly192 Accession # Q8CD15
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS.

## APPLICATIONS

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
<b>Western Blot</b>	2 µg/mL	See Below

## DATA



## PREPARATION AND STORAGE

<b>Reconstitution</b>	Sterile PBS to a final concentration of 0.5 mg/mL.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
<b>Stability &amp; Storage</b>	<b>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</b> <ul style="list-style-type: none"> <li>• 12 months from date of receipt, -20 to -70 °C as supplied.</li> <li>• 1 month, 2 to 8 °C under sterile conditions after reconstitution.</li> <li>• 6 months, -20 to -70 °C under sterile conditions after reconstitution.</li> </ul>

## BACKGROUND

MINA (myc-induced nuclear antigen; also Mina53) is a 52-54 kDa member of both the MINA53/NO66 and Jumonji C family of proteins. Its expression is associated with proliferating cells, and it has been found in cytoplasm, nucleus and nucleoli. MINA appears to be induced by c-myc, and synthesized by spermatogonia, occasional squamous epithelium, naïve T cells and select cancer cells. When expressed, MINA is reported to regulate expression of genes such as HGF, EGF-R and IL-4. It may exert its regulatory activity through an intrinsic demethylase function. Mouse MINA is 465 amino acids (aa) in length. It possesses one cupin (or enzyme-associated) region (aa 51-363) that contains a JmjC domain (aa 139-271). There are two potential isoform variants that contain either a 12 aa substitution for aa 145-465, or a 15 aa substitution for aa 228-465. Over aa 2-192, mouse MINA shares 92% and 82% aa sequence identity with rat and human MINA, respectively.