

StemXVivo[®]

Cardiomyocyte Differentiation Kit

Catalog # SC032B

StemXVivo[®]

Cardiomyocyte Differentiation Kit

Catalog Number SC032B

Reagents for the directed differentiation of human pluripotent stem cells into the cardiomyocyte lineage.

This package insert must be read in its entirety before using this product.
For laboratory research use only. Not for diagnostic use.
The safety and efficacy of this product in diagnostic or
other clinical uses has not been established.

TABLE OF CONTENTS

SECTION	PAGE
INTRODUCTION	1
PRINCIPLE OF THE ASSAY.....	1
TECHNICAL HINTS & LIMITATIONS	2
PRECAUTION	2
MATERIALS PROVIDED & STORAGE CONDITIONS	3
OTHER SUPPLIES REQUIRED	4
REAGENT & MATERIAL PREPARATION.....	5
PROCEDURE OUTLINE	6
UNDIFFERENTIATED CELL PREPARATION.....	7
CARDIOMYOCYTE DIFFERENTIATION PROCEDURE.....	8
FIXING & STAINING PROCEDURE	10
DATA EXAMPLES	11
REFERENCES	11

MANUFACTURED AND DISTRIBUTED BY:

USA & Canada | R&D Systems, Inc.

614 McKinley Place NE, Minneapolis, MN 55413, USA
TEL: (800) 343-7475 (612) 379-2956 FAX: (612) 656-4400
E-MAIL: info@RnDSystems.com

DISTRIBUTED BY:

UK & Europe | R&D Systems Europe, Ltd.

19 Barton Lane, Abingdon Science Park, Abingdon OX14 3NB, UK
TEL: +44 (0)1235 529449 FAX: +44 (0)1235 533420
E-MAIL: info@RnDSystems.co.uk

China | R&D Systems China Co., Ltd.

24A1 Hua Min Empire Plaza, 726 West Yan An Road, Shanghai PRC 200050
TEL: +86 (21) 52380373 FAX: +86 (21) 52371001
E-MAIL: info@RnDSystemsChina.com.cn

INTRODUCTION

Pluripotent stem cells, including both embryonic and induced pluripotent stem cells, provide much promise for the generation of sufficient quantities of specialized cells for use in regenerative medicine. Additionally, these cells are an important tool for understanding developmental and disease mechanisms.

The ability of pluripotent stem cells to differentiate into functional cardiomyocytes holds tremendous promise for applications in therapeutics, pharmaceuticals, and developmental biology research. The potentially unlimited supply of cardiomyocytes generated from pluripotent cell sources can be used in high throughput platforms for drug toxicity screening, eliminating harmful drug candidates before clinical trial stages. Additionally, fully characterized differentiated cardiomyocytes may eventually be transplanted as a therapy for heart disease. Studying the process of cardiomyocyte differentiation from pluripotent cells will increase our understanding of the processes involved in embryonic development and organogenesis (1-3).

PRINCIPLE OF THE ASSAY

The StemXVivo® Cardiomyocyte Differentiation Kit contains specially formulated media supplements and growth factors that can be used to direct differentiation of pluripotent stem cells toward a cardiomyocyte fate. An antibody is included to characterize the extent of differentiation. The quantity of each component in the kit is sufficient to differentiate two 24-well plates, or an equivalent surface area, of pluripotent stem cells.

TECHNICAL HINTS & LIMITATIONS

- FOR LABORATORY RESEARCH USE ONLY. NOT FOR DIAGNOSTIC USE.
- Do not mix or substitute reagents with those from other lots or sources.
- The safety and efficacy of this product in diagnostic or other clinical uses has not been established.
- The quality and differentiation potential of human pluripotent stem cells at the onset of the differentiation protocol are of paramount importance to the efficiency of differentiation.
- If no beating is observed during the differentiation, test the cells' ability to differentiate into mesoderm by either differentiating with the StemXVivo® Mesoderm Kit (R&D Systems, Catalog # SC030B) or by staining the cells with Goat Anti-Human/Mouse Brachyury Affinity-purified Polyclonal Antibody (R&D Systems, Catalog # AF2085) after the first 24 hours of differentiation (prior to the Day 1 Media Change).
- If beating is observed, but with low efficiency, the quality of the starting population should be verified at Day 0 of the differentiation (prior to the Day 0 Media Change). Cell quality can be determined morphologically, by staining with pluripotency markers such as Oct-4A (R&D Systems, Catalog # MAB17591), Nanog (R&D Systems, Catalog # AF1997), SSEA-4 (R&D Systems, Catalog # MAB1435), or others, or functionally by using the Human Pluripotent Stem Cell Functional Identification Kit (R&D Systems, Catalog # SC027B) to determine the ability of the starting cell population to differentiate into each of the three germ layers.

PRECAUTION

The acute and chronic effects of overexposure to reagents of this kit are unknown. Safe laboratory procedures should be followed and protective clothing should be worn when handling kit reagents.

MATERIALS PROVIDED & STORAGE CONDITIONS

Store unopened kit at $\leq -70\text{ }^{\circ}\text{C}$ in a manual defrost freezer. Do not use past kit expiration date.

PART	PART #	DESCRIPTION	STORAGE OF OPENED/ RECONSTITUTED MATERIAL
Stem Cell Qualified RGF BME, Pathclear®	894860	1.5 mL of frozen reduced growth factor basement membrane extract.	Store at 2-8 °C on ice for up to 1 week or aliquot and store at $\leq -70\text{ }^{\circ}\text{C}$ for up to 3 months.* Avoid repeated freeze-thaw cycles.
Cardiomyocyte Differentiation Base Media Supplement I	390537	5.5 mL of a 50X concentrated solution.	Store at 2-8 °C for up to 2 weeks or aliquot and store at $\leq -20\text{ }^{\circ}\text{C}$ in a manual defrost freezer for up to 6 months.* Avoid repeated freeze-thaw cycles.
Cardiomyocyte Differentiation Base Media Supplement II	390538	10 mL of a 50X concentrated solution.	
Cardiomyocyte Differentiation Cocktail I	967589	1 vial of lyophilized growth factors; enough to make 25 μL of a 1000X stock.	Store under sterile conditions at 2-8 °C for up to 1 month or aliquot and store at $\leq -20\text{ }^{\circ}\text{C}$ in a manual defrost freezer for up to 6 months.* Avoid repeated freeze-thaw cycles.
Cardiomyocyte Differentiation Cocktail IIA	390600	1 vial of lyophilized growth factors; enough to make 50 μL of a 1000X stock.	
Cardiomyocyte Differentiation Cocktail IIB	390601	1 vial containing 10 μL of a 5000X stock solution	
Cardiomyocyte Differentiation Cocktail III	967590	1 vial of lyophilized growth factors; enough to make 50 μL of a 1000X stock.	
Anti-Human Cardiac Troponin T	893409	25 μg of lyophilized mouse anti-human Troponin T monoclonal antibody. Enough to make 2.5 mL of staining solution when used at the suggested concentration of 10 $\mu\text{g}/\text{mL}$.	

*Provided this is within the expiration of the kit.

OTHER SUPPLIES REQUIRED

Materials

- Human pluripotent stem cells
- 24-well culture plates (or other, as needed)
- 60 mm culture plates
- 12 mm coverslips (optional)
- 15 mL and 50 mL centrifuge tubes
- 0.2 μ m syringe filter
- 10 mL syringe
- Pipettes and pipette tips
- Serological pipettes
- Glass slides
- Fine pointed curved forceps
- FACS tubes

Reagents

- Pluripotent Stem Cell Maintenance Media (R&D Systems, Catalog # AR005) or equivalent
- Cultrex® Stem Cell Qualified Reduced Growth Factor Base Membrane Extract, Pathclear® (R&D Systems, Catalog # 3434-001-02)
- RPMI 1640
- BSA, very low endotoxin
- DMEM/F-12 (1X)
- GlutaMAX™ (Invitrogen, Catalog # 35050-079 or equivalent)
- Penicillin-Streptomycin (optional)
- Phosphate Buffered Saline (PBS)
- ROCK Inhibitor (Y-27632; Tocris®, Catalog # 1254)
- Accutase® or Versene™
- 95% Ethanol
- 4% Paraformaldehyde
- 1% BSA in PBS
- 0.3% Triton X-100, 1% BSA, 10% normal donkey serum in PBS
- Mounting Medium (R&D Systems, Catalog # CTS011)
- Secondary antibody for immunocytochemistry (R&D Systems, Catalog # NL007)
- Deionized or distilled water

Equipment

- 37°C and 5% CO₂ incubator
- Centrifuge
- Inverted microscope
- Fluorescence microscope
- Hemocytometer

REAGENT & MATERIAL PREPARATION

Cultrex® Stem Cell Qualified Reduced Growth Factor Basement Membrane Extract, Pathclear® (RGF BME) - Store tubes of RGF BME at ≤ -70 °C until ready to use. As needed, thaw RGF BME overnight on ice at 2-8 °C. Dilute 1:40 in ice cold Pluripotent Stem Cell Maintenance Media for coating the tissue culture dish prior to seeding cells for differentiation. Dilute 1:60 in ice cold Pluripotent Stem Cell Maintenance Media for overlay procedure on Day (-1). Dilute 1:60 in Cardiomyocyte Differentiation Base Media I to initiate differentiation on Day 0.

0.1% BSA in PBS - Dissolve 10 mg of BSA in 10 mL of PBS. Sterile filter the solution by syringe filter and store at 2-8 °C for up to 3 months.

Cardiomyocyte Differentiation Cocktail I (1000X) - Reconstitute with 25 μ L of sterile 0.1% BSA in PBS. Mix gently.

Cardiomyocyte Differentiation Cocktail IIA (1000X) - Reconstitute with 50 μ L of sterile 0.1% BSA in PBS. Mix gently.

Cardiomyocyte Differentiation Cocktail III (1000X) - Reconstitute with 50 μ L of sterile 0.1% BSA in PBS. Mix gently.

Cardiomyocyte Differentiation Base Media I - Dilute Cardiomyocyte Differentiation Base Media Supplement I 1:50 with RPMI 1640 containing 1X Penicillin/Streptomycin (optional) and 1X GlutaMAX. Prepare fresh as needed.

Note: *On Day 0, the media should be prepared on ice immediately prior to use.*

Cardiomyocyte Differentiation Base Media II - Dilute Cardiomyocyte Differentiation Base Media Supplement II 1:50 with RPMI 1640 containing 1X Penicillin/Streptomycin (optional) and 1X GlutaMAX. Prepare fresh as needed.

PROCEDURE OUTLINE

Coat wells with Stem Cell Qualified PathClear RGF BME (RGF BME).
Incubate at room temperature for 1-2 hours.







Plate human pluripotent stem cells onto the coated plates at $3-4 \times 10^4$ cells/cm² in Pluripotent Stem Cell Maintenance Media.
Culture cells to 80-90% confluency.




Day (-1) of Differentiation
Replace the stem cell culture media with ice cold Pluripotent Stem Cell Maintenance Media containing RGF BME diluted 1:60.
Incubate at 37 °C and 5% CO₂ for 18-24 hours.




Day 5 of Differentiation
Replace the media with Day 5 Cardiomyocyte Differentiation Media.
Incubate at 37 °C and 5% CO₂ for 2 DAYS without media exchange.




Day 0 of Differentiation
Replace the media with ice cold Day 0 Cardiomyocyte Differentiation Media containing RGF BME diluted 1:60.
Incubate at 37 °C and 5% CO₂ for 24 hours.




Day 7 of Differentiation and Beyond
Replace the media with Cardiomyocyte Differentiation Base Media I.
Incubate at 37 °C and 5% CO₂. Replace media every 1-2 days as needed.



Day 1 of Differentiation
Replace the media with Day 1 Cardiomyocyte Differentiation Media.
Incubate at 37 °C and 5% CO₂ for 4 DAYS without media exchange.



Day 12 of Differentiation and Beyond
Replace the media with Cardiomyocyte Differentiation Base Media II.
Incubate at 37 °C and 5% CO₂. Replace media every 1-2 days as needed.



UNDIFFERENTIATED CELL PREPARATION

Note: *This protocol has been tested on cells cultured in either MEF Conditioned Media (R&D Systems, Catalog # AR005) or equivalent.*

The quality and differentiation potential of human pluripotent stem cells at the onset of the differentiation protocol are of paramount importance to the efficiency of differentiation. Human pluripotent stem cells must be > 95% positive for OCT-3/4.

COATING PLATES WITH RGF BME

1. Thaw Cultrex® Stem Cell Qualified RGF BME, Pathclear® (RGF BME) on ice at 2-8 °C overnight.
2. Dilute the RGF BME 1:40 in ice cold DMEM/F-12. This can be stored at 2-8 °C for up to 2 weeks.
3. Coat the desired number of wells or plates with diluted RGF BME (0.5 mL/well for a 24-well plate) and incubate for 1-2 hours at room temperature.
4. RGF BME can be kept on ice at 2-8 °C for up to 2 weeks. Avoid repeated freeze/thaw cycles.

CELL DISSOCIATION

1. Warm the pluripotent stem cell maintenance media to room temperature.
2. Remove the existing media from the cells. Add 1 mL of Accutase or Versene solution to each 60 mm plate or 3 mL to each 100 mm plate. Incubate at room temperature for 2-7 minutes or until the cells begin to slough off the plate. If using cells from several plates, work in small batches (1-2 plates at a time) so the cells are not exposed to the Accutase or Versene beyond the time it takes the cells to slough off the plate.
3. Pipette gently over the plate until the cells become detached.
4. Gently pipette the cell suspension up and down to break up large cell clumps.
5. Transfer the cell suspension to a 15 mL centrifuge tube containing 5 mL of pluripotent stem cell maintenance media (or 12 mL if using a 100 mm plate) and spin at 200 x g for 4 minutes.

CELL PLATING

1. Resuspend the pellet in pluripotent stem cell maintenance media, and count the viable cells using Trypan Blue and a hemocytometer.
2. Plate the cells onto prepared RGF BME-coated plates at a concentration of $3.5-4 \times 10^4$ cells/ cm². It is recommended to add the ROCK inhibitor (Y-27632) for the first 16-24h of culture.
3. Allow the cells to attach and grow at 37 °C and 5% CO₂ with daily media exchanges until the monolayer reaches approximately 80-90% confluence (3-4 days).

CARDIOMYOCYTE DIFFERENTIATION PROCEDURE

RGF BME Overlay: Day (-1) through Day 0

1. Pluripotent cells should be grown to approximately 80-90% confluence on RGF BME prior to the initiation of differentiation.
2. For this step, Pluripotent Stem Cell Maintenance Media should be kept on ice and used ice cold.
3. One tube of RGF BME should be thawed overnight and ready for dilution (see Reagent and Material Preparation on Page 5).
4. Quickly dilute an appropriate amount of RGF BME 1:60 in ice cold Pluripotent Stem Cell Maintenance Media. For 24-well plates, use 0.5 mL/well.
5. Quickly remove and discard spent media and add the RGF BME-containing media.
Note: Steps 4 and 5 should be completed as quickly as possible, without sacrificing sterile technique, to prevent the RGF BME from starting to gel into solid clumps in the media.
6. Culture the cells overnight at 37 °C and 5% CO₂.
7. RGF BME can be kept on ice at 2-8 °C for up to 2 weeks. Avoid repeated freeze/thaw cycles.

RGF BME Overlay and Differentiation Initiation: Day 0 through Day 1

1. Prepare Day 0 Cardiomyocyte Differentiation Media as follows:
 - a. Determine beforehand the appropriate total volume of media needed. For 24-well plates, use 0.5 mL/well.
 - b. Pipette the needed volume of ice cold Cardiomyocyte Differentiation Base Media I into a sterile centrifuge tube.
 - c. Quickly dilute RGF BME 1:60 into the ice cold Media.
 - d. Quickly dilute the Cardiomyocyte Differentiation Cocktail I 1:1000 into the ice cold Media.
 - e. Mix gently.
2. Remove and discard the spent media and add the Day 0 Cardiomyocyte Differentiation Media to each well.
Note: Steps 1 and 2 should be completed as quickly as possible, without sacrificing sterile technique, to prevent the RGF BME from starting to gel into solid clumps in the media.
3. Culture the cells overnight at 37 °C and 5% CO₂.

CARDIOMYOCYTE DIFFERENTIATION PROCEDURE *CONTINUED*

Differentiation Days 1 through 5

1. Prepare Day 1 Cardiomyocyte Differentiation Media as follows:
 - a. Warm the appropriate volume of Cardiomyocyte Differentiation Base Media I to room temperature. For 24-well plates, use 1 mL/well.
 - b. Dilute the Cardiomyocyte Differentiation Cocktail IIA 1:1000 and Cardiomyocyte Differentiation Cocktail IIB 1:5000 into the Base Media I.
 - c. Mix gently.
2. Remove and discard the spent media and add the Day 1 Cardiomyocyte Differentiation Media to each well.
3. Culture the cells for four days (Day 1-5) at 37 °C and 5% CO₂ without media exchange.

Note: *The media will turn very yellow as the pH increases, but it is imperative to not remove the media.*

Differentiation Days 5 though 7

1. Prepare Day 5 Cardiomyocyte Differentiation Media as follows:
 - a. Warm the appropriate volume of Cardiomyocyte Differentiation Base Media I to room temperature. For 24-well plates, use 1 mL/well.
 - b. Dilute the Cardiomyocyte Differentiation Cocktail III 1:1000 into the Base Media I.
 - c. Mix gently.
2. Remove and discard the spent media and add the Day 5 Cardiomyocyte Differentiation Media to each well.
3. Culture the cells for two days (Day 5-7) at 37 °C and 5% CO₂ without media exchange.

Differentiation Day 7 through 12

1. Warm the Cardiomyocyte Differentiation Base Media I to room temperature.
2. Remove and discard the spent media and add the prewarmed Cardiomyocyte Differentiation Base Media I to each well. For 24-well plates, use 1 mL/well.
3. Culture the cells at 37 °C and 5% CO₂. Exchange the media every one to two days.

Note: *Initially, media exchange every second day is standard. As beating cells begin to emerge, it may become necessary to exchange the media every day.*

Differentiation Day 12 and Beyond

1. Warm the Cardiomyocyte Differentiation Base Media II to room temperature.
2. Remove and discard the spent media and add the prewarmed Cardiomyocyte Differentiation Base Media II to each well. For 24-well plates, use 1 mL/well.
3. Culture the cells at 37 °C and 5% CO₂. Exchange the media every one to two days.

FIXING & STAINING PROCEDURE

Immunocytochemistry

1. Wash the cells twice with PBS (1 mL/well for a 24-well plate).
2. Fix the cells with 4% paraformaldehyde in PBS for 20 minutes at room temperature.
3. Wash the cells 3 times with 1% BSA in PBS for 5 minutes (0.5 mL/well for a 24-well plate).
4. Permeabilize and block the cells with 0.3% Triton X-100, 1% BSA, and 10% normal donkey serum in PBS at room temperature for 45 minutes (0.5 mL/well for a 24-well plate).
5. During the blocking, dilute the reconstituted Anti-Human Cardiac Troponin T Primary Antibody to a final concentration of 10 ug/mL in PBS containing 0.3% Triton X-100, 1% BSA, and 10% normal donkey serum.

Note: A negative control should be performed using PBS containing 0.3% Triton X-100, 1% BSA, and 10% normal donkey serum in the absence of a primary antibody.

6. After blocking, incubate the cells with diluted Anti-Human Cardiac Troponin T Primary Antibody (300 µL/well of a 24-well plate) for 3 hours at room temperature or overnight at 2-8 °C.
7. Wash the cells 3 times with 1% BSA in PBS for 5 minutes (0.5 mL/well of a 24-well plate).
8. Dilute the secondary antibody [e.g., NL557-conjugated Donkey Anti-Mouse Secondary Antibody (R&D Systems, Catalog #NL007)] at 1:200 in PBS containing 1% BSA.
9. Incubate the cells with diluted secondary antibody in the dark for 60 minutes at room temperature (300 µL/well of a 24-well plate).
10. Wash the cells 3 times with 1% BSA in PBS for 5 minutes (0.5 mL/well of a 24-well plate).
11. Cover the cells with PBS (1 mL/well of a 24-well plate) and visualize with a fluorescence microscope. Alternatively, aspirate the PBS and add distilled or deionized water (0.5 mL/well of a 24-well plate). Carefully remove each coverslip with forceps and mount cell-side down onto a drop of mounting medium on a glass slide.
12. Slides are ready for microscopic observation.

DATA EXAMPLES




Figure 1: Differentiation of Pluripotent Stem Cells into Cardiomyocytes. JOY6 (left panel) and iBJ6 (right panel) human induced pluripotent stem cells were differentiated into cardiomyocytes using the media supplements included in this kit. Aside from visually observing contracting cells, commitment to the cardiomyocyte cell fate was evaluated by labeling with the Anti-Human Cardiac Troponin T antibody included in this kit. For visualization, the cells were stained using NorthernLights™ 557-conjugated Donkey anti-Mouse Secondary Antibody (R&D Systems, Catalog # NL007; red), and the nuclei were counterstained with DAPI (blue).

REFERENCES

1. Dambrot, C. *et al.* (2011) *Biochem. J.* **434**: 25.
2. Laflamme, M.A. *et al.* (2007) *Nat. Biotech.* **25**: 1015.
3. Xu C. *et al.* (2002) *Circ. Res.* **91**: 501.

NOTES

NOTES

NOTES

*BG01V human embryonic stem cells are licensed from Viacyste, Inc.
All trademarks and registered trademarks are the property of their respective owners.*

©2016 R&D Systems, Inc.

10.16

727005.0

10/16