

## MEF2a(Phospho-Thr312) Antibody

Catalog No: #11039

Package Size: #11039-1 50ul #11039-2 100ul #11039-4 25ul

Orders: order@signalwayantibody.com

Support: tech@signalwayantibody.com

## Description

|                       |   |
|-----------------------|---|
| Product Name          | MEF2a(Phospho-Thr312) Antibody  |
| Host Species          | Rabbit  |
| Clonality             | Polyclonal  |
| Purification          | Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates.<br>Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatography using non-phosphopeptide. |
| Applications          | WB IHC IF   |
| Species Reactivity    | Hu Ms Rt  |
| Specificity           | The antibody detects endogenous level of MEF2A only when phosphorylated at threonine 312.   |
| Immunogen Type        | Peptide-KLH   |
| Immunogen Description | Peptide sequence around phosphorylation site of threonine 312 (L-A-T(p)-P-V) derived from Human MEF2A.  |
| Target Name           | MEF2a   |
| Modification          | Phospho-Thr312  |
| Other Names           | MEF2, ADCAD1, RSRFC4, RSRFC9  |
| Accession No.         | Swiss-Prot: Q02078NCBI Protein: NP_001124398.1  |
| Concentration         | 1.0mg/ml  |
| Formulation           | Supplied at 1.0mg/mL in phosphate buffered saline (without Mg <sup>2+</sup> and Ca <sup>2+</sup> ), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.  |
| Storage               | Store at -20°C for long term preservation (recommended). Store at 4°C for short term use.   |


## Application Details

Predicted MW: 54kd


Immunohistochemistry: 1:50~1:100

Immunofluorescence: 1:100~1:200


## Images



Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using MEF2A(Phospho-Thr312) Antibody #11039(left) or the same antibody preincubated with blocking peptide(right).



Immunofluorescence staining of methanol-fixed HeLa cells using MEF2A(Phospho-Thr312) Antibody #11039.



Western blot analysis of extracts from C2C12 cells, treated with Anisomycin or calf intestinal phosphatase (CIP), using MEF2A (Phospho-Thr312) Antibody #11039.

## Background

The process of differentiation from mesodermal precursor cells to myoblasts has led to the discovery of a variety of tissue-specific factors that regulate muscle gene expression. The myogenic basic helix-loop-helix proteins, including myoD (MIM 159970), myogenin (MIM 159980), MYF5 (MIM 159990), and MRF4 (MIM 159991) are one class of identified factors. A second family of DNA binding regulatory proteins is the myocyte-specific enhancer factor-2 (MEF2) family. Each of these proteins binds to the MEF2 target DNA sequence present in the regulatory regions of many, if not all, muscle-specific genes. The MEF2 genes are members of the MADS gene family (named for the yeast mating type-specific transcription factor MCM1, the plant homeotic genes 'agamous' and 'deficiens' and the human serum response factor SRF (MIM 600589)), a family that also includes several homeotic genes and other transcription factors, all of which share a conserved DNA-binding domain

K Satoh, J Ohnishi, A Sato, et al. (2007) Nemo-Like Kinase-Myocyte Enhancer Factor 2A Signaling Regulates Anterior Formation in Xenopus Development. *Molecular and Cellular Biology*, 27(21):7623-30.

This article references the use of the #11039 in the following applications :Western blotting

## Published Papers

K Satoh, J Ohnishi, A Sato et al., Nemo-Like Kinase-Myocyte Enhancer Factor 2A Signaling Regulates Anterior Formation in Xenopus Development., *Molecular and Cellular Biology*, 27(21):7623-30(2007)

[PMID:17785444](https://pubmed.ncbi.nlm.nih.gov/17785444/)

Note: This product is for in vitro research use only and is not intended for use in humans or animals.