## ATF4(Phospho-Ser245) Antibody

Catalog No: #11053

Package Size: #11053-1 50ul #11053-2 100ul #11053-4 25ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

	$\alpha$	∩rı	n	t١	$\smallfrown$	n
u	es		u	ш	u	ш
			ш.			


Product Name	ATF4(Phospho-Ser245) Antibody	
Host Species	Rabbit	
Clonality	Polyclonal	
Purification	Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates.	
	Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho	
	specific antibodies were removed by chromatogramphy using non-phosphopeptide.	
Applications	IHC WB	
Species Reactivity	Hu	
Specificity	The antibody detects endogenous level of ATF4 only when phosphorylated at serine 245.	
Immunogen Type	Peptide-KLH	
Immunogen Description	Peptide sequence around phosphorylation site of serine 245 (N-R-S(p)-L-P) derived from Human ATF4.	
Target Name	ATF4	
Modification	Phospho-Ser245	
Other Names	C/ATF; C/EBP-related ATF; CREB2	
Accession No.	Swiss-Prot: P18848NCBI Protein: NP _001666.2	
Concentration	1.0mg/ml	
Formulation	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02%	
	sodium azide and 50% glycerol.	
Storage	Store at -20°C for long term preservation (recommended). Store at 4°C for short term use.	

## **Application Details**


Predicted MW: 45kd

Western blotting: 1:500~1:1000
Immunohistochemistry: 1:50~1:100

## **Images**



Western blot analysis of extracts from Hela cells untreated or treated with UV using ATF4(Phospho-Ser245) Antibody #11053.



Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using ATF4(Phospho-Ser245) Antibody #11053(left) or the same antibody preincubated with blocking peptide(right).

## Background

ATF4 encodes a transcription factor that was originally identified as a widely expressed mammalian DNA binding protein that could bind a tax-responsive enhancer element in the LTR of HTLV-1. The encoded protein was also isolated and characterized as the cAMP-response element binding protein 2 (CREB-2). The protein encoded by this gene belongs to a family of DNA-binding proteins that includes the AP-1 family of transcription factors, cAMP-response element binding proteins (CREBs) and CREB-like proteins. These transcription factors share a leucine zipper region that is involved in protein-protein interactions, located C-terminal to a stretch of basic amino acids that functions as a DNA binding domain. Two alternative transcripts encoding the same protein have been described. Two pseudogenes are located on the X chromsome at q28 in a region containing a large inverted duplication.

Yang X, et al. (2004). Cell.117(3): 387-398.

Note: This product is for in vitro research use only and is not intended for use in humans or animals.