HER2(Phospho-Tyr1221/Tyr1222) Antibody

Catalog No: #11076

Description

Package Size: #11076-1 50ul #11076-2 100ul #11076-4 25ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

| Product Name | HER2(Phospho-Tyr1221/Tyr1222) Antibody |
|-----------------------|--|
| Host Species | Rabbit |
| Clonality | Polyclonal |
| Purification | Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. |
| | Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho |
| | specific antibodies were removed by chromatogramphy using non-phosphopeptide. |
| Applications | WB IF |
| Species Reactivity | Human |
| Specificity | The antibody detects endogenous level of HER2 only when phosphorylated at tyrosine1221/tyrosine1222. |
| Immunogen Type | Peptide-KLH |
| Immunogen Description | Peptide sequence around phosphorylation site of tyrosine1221/1222 (N-L-Y(p)-Y(p)-W) derived from Human |
| | HER2. |
| Target Name | HER2 |
| Modification | Phospho-Tyr1221/Tyr1222 |
| Other Names | C-erbB-2; ErbB2; |
| | |

Swiss-Prot: P04626NCBI Protein: NP_001005862.1

sodium azide and 50% glycerol.

Application Details

Accession No.

Concentration

Formulation


Storage

Predicted MW: 185kd

Western blotting: 1:500~1:1000

Immunofluorescence: 1:100~1:200


Images




Western blot analysis of extracts from MDA cells untreated or treated with EGF using HER2(Phospho-Tyr1221/Tyr1222) Antibody #11076.

Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02%

Store at -20°C for long term preservation (recommended). Store at 4°C for short term use.



Immunofluorescence staining of methanol-fixed MCF cells using HER2(Phospho-Tyr1221/Tyr1222) Antibody #11076.



Western blot analysis of extracts from Hela cells, treated with EGF or calf intestinal phosphatase (CIP), using HER2 (Phospho-Tyr1221/Tyr1222) Antibody #11076.

Background

Essential component of a neuregulin-receptor complex, although neuregulins do not interact with it alone. GP30 is a potential ligand for this receptor. Not activated by EGF, TGF-a and amphiregulin.

Marone R, et al. (2004) Nat Cell Biol; 6(6): 515-22.

Ren Z, et al. (2002) J Biol Chem; 277(41): 38486-93.

Note: This product is for in vitro research use only and is not intended for use in humans or animals.