

## VEGFR2(Phospho-Tyr1175) Antibody

Catalog No: #11084

Package Size: #11084-1 50ul #11084-2 100ul #11084-4 25ul

Orders: [order@signalwayantibody.com](mailto:order@signalwayantibody.com)Support: [tech@signalwayantibody.com](mailto:tech@signalwayantibody.com)

## Description

Product Name	VEGFR2(Phospho-Tyr1175) Antibody
Host Species	Rabbit
Clonality	Polyclonal
Purification	Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatography using non-phosphopeptide.
Applications	IHC IF
Species Reactivity	Hu Ms Rt
Specificity	The Antibody detects endogenous level of VEGFR2 only when phosphorylated at tyrosine 1175.
Immunogen Type	Peptide-KLH
Immunogen Description	Peptide sequence around phosphorylation site of tyrosine 1175 (K-D-Y(p)-I-V) derived from Human VEGFR2.
Target Name	VEGFR2
Modification	Phospho-Tyr1175
Other Names	FLK1; KDR; VGFR2; VGR2; kinase insert domain receptor
Accession No.	Swiss-Prot: P35968NCBI Protein: NP_002244.1
Concentration	1.0mg/ml
Formulation	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg <sup>2+</sup> and Ca <sup>2+</sup> ), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Storage	Store at -20°C for long term preservation (recommended). Store at 4°C for short term use.


## Application Details

Predicted MW: 230kd


Immunohistochemistry: 1:50~1:100

Immunofluorescence: 1:100~1:200

## Images



Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using VEGFR2(Phospho-Tyr1175) Antibody #11084(left) or the same antibody preincubated with blocking peptide(right).



Immunofluorescence staining of methanol-fixed MCF cells using VEGFR2(Phospho-Tyr1175) Antibody #11084.

## Background

Receptor for VEGF or VEGFC. Has a tyrosine-protein kinase activity. The VEGF-kinase ligand/receptor signaling system plays a key role in vascular development and regulation of vascular permeability. In case of HIV-1 infection, the interaction with extracellular viral Tat protein seems to enhance angiogenesis in Kaposi's sarcoma lesions

Lamalice L, et al. (2004). *Oncogene*.23(2): 434-445.

Takahashi T, et al. (2001). *EMBO J* .20(11): 2768-2778.

Note: This product is for in vitro research use only and is not intended for use in humans or animals.