b-Catenin(Phospho-Ser33) Antibody

Catalog No: #11218

Package Size: #11218-1 50ul #11218-2 100ul #11218-4 25ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

| Descr | iption |
|-------|--------|
| D | N1 |

| Product Name | b-Catenin(Phospho-Ser33) Antibody |
|-----------------------|--|
| Host Species | Rabbit |
| Clonality | Polyclonal |
| Purification | Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. |
| | Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho |
| | specific antibodies were removed by chromatogramphy using non-phosphopeptide. |
| Applications | WB IHC IF |
| Species Reactivity | Hu Ms Rt |
| Specificity | The antibody detects endogenous level of b-Catenin only when phosphorylated at serine 33. |
| Immunogen Type | Peptide-KLH |
| Immunogen Description | Peptide sequence around phosphorylation site of serine 33 (L-D-S(p)-G-I) derived from Human b-Catenin. |
| Target Name | b-Catenin |
| Modification | Phospho-Ser33 |
| Other Names | CTNNB1; CATNB; CTNB1; CTNNB; |
| Accession No. | Swiss-Prot: P35222NCBI Protein: NP_001091679.1 |
| Concentration | 1.0mg/ml |
| Formulation | Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% |
| | sodium azide and 50% glycerol. |
| Storage | Store at -20°C for long term preservation (recommended). Store at 4°C for short term use. |
| | |


Application Details

Predicted MW: 92kd


Western blotting: 1:500~1:1000

Immunohistochemistry: 1:50~1:100

Images



Western blot analysis of extracts from MCF-7 cells untreated or treated with Calyculin A using b-Catenin(Phospho-Ser33) Antibody #11218.



Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using b-Catenin(Phospho-Ser33) Antibody #11218(left) or the same antibody preincubated with blocking peptide(right).

Background

Involved in the regulation of cell adhesion and in signal transduction through the Wnt pathway.

Novak A, et al. (1998) Proc Natl Acad Sci U S A; 95(8): 4374-4379

Marin O, et al. (2003) Proc Natl Acad Sci U S A; 100(18): 10193-10200

Okamura H, et al. (2004) Mol Cell Biol; 24(10): 4184-4195

Xing Y, et al. (2003) Genes Dev; 17(22): 2753-2764

Barth AI, et al. (1999) Proc Natl Acad Sci U S A; 96(9): 4947-4952

Note: This product is for in vitro research use only and is not intended for use in humans or animals.