Product Datasheet

PKR(Phospho-Thr451) Antibody

Catalog No: #11290

Package Size: #11290-1 50ul #11290-2 100ul #11290-4 25ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Description Product Name PKR(Phospho-Thr451) Antibody Host Species Rabbit


| Product Name | PKR(Phospho-Thr451) Antibody |
|-----------------------|--|
| Host Species | Rabbit |
| Clonality | Polyclonal |
| Purification | Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. |
| | Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho |
| | specific antibodies were removed by chromatogramphy using non-phosphopeptide. |
| Applications | IHC |
| Species Reactivity | Hu |
| Specificity | The antibody detects endogenous level of PKR only when phosphorylated at threonine 451. |
| Immunogen Type | Peptide-KLH |
| Immunogen Description | Peptide sequence around phosphorylation site of threonine 451 (K-G-T(p)-L-R) derived from Human PKR. |
| Target Name | PKR |
| Modification | Phospho-Thr451 |
| Other Names | ADRB2; E2AK2; EIF2AK2; EIF2aK; PRKR |
| Accession No. | Swiss-Prot: P19525 NCBI Protein: NP_001129123.1 |
| Concentration | 1.0mg/ml |
| Formulation | Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% |
| | sodium azide and 50% glycerol. |
| Storage | Store at -20°C for long term preservation (recommended). Store at 4°C for short term use. |

Application Details

Predicted MW: 68kd

Immunohistochemistry: 1:50~1:100

Images



Immunohistochemical analysis of paraffin-embedded human colon carcinoma tissue using PKR(Phospho-Thr451) Antibody #11290(left) or the same antibody preincubated with blocking peptide(right).

Background

Following activation by double-stranded RNA in the presence of ATP, the kinase becomes autophosphorylated and can catalyze the phosphorylation of the translation initiation factor EIF2S1, which leads to an inhibition of the initiation of protein synthesis. Double-stranded RNA is generated during the course of a viral infection.

Ingrid K. Ruf, et,al. (2005) J. Virol ; 79: 14562 - 14569.

Christy M. Hebner, et, al. (2006) J. Gen. Virol ; 87: 3183 - 3193.

Rika van Huizen, et,al. (2003) J. Biol. Chem ; 278: 15558 - 15564.

Note: This product is for in vitro research use only and is not intended for use in humans or animals.