PKCd(Phospho-Ser645) Antibody

Catalog No: #11296

Package Size: #11296-1 50ul #11296-2 100ul #11296-4 25ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Description

Product Name	PKCd(Phospho-Ser645) Antibody		
Host Species	Rabbit		
Clonality	Polyclonal		
Purification	Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates.		
	Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho		
	specific antibodies were removed by chromatogramphy using non-phosphopeptide.		
Applications	WB IHC		
Species Reactivity	Hu Rt		
Specificity	The antibody detects endogenous level of PKCd only when phosphorylated at serine 645.		
Immunogen Type	Peptide-KLH		
Immunogen Description	Peptide sequence around phosphorylation site of serine 645 (R-L-S(p)-Y-S) derived from Human PKCd.		
Target Name	PKCd		
Modification	Phospho-Ser645		
Other Names	KPCD; PKC-delta; PRKCD; kinase PKC-delta; nPKC-delta		
Accession No.	Swiss-Prot: Q05655NCBI Protein: NP_006245.2		
Concentration	1.0mg/ml		
Formulation	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02%		
	sodium azide and 50% glycerol.		
Storage	Store at -20°C for long term preservation (recommended). Store at 4°C for short term use.		


Application Details

Predicted MW: 78kd			
Western blotting: 1:500~1:1000			
Immunohistochemistry: 1:50~1:	00		

Images

КD 95 —	PC12
72 —	PKCð (pSer645)
55 —	
43 — PMA	- +

Western blot analysis of extracts from PC12 cells untreated or treated with PMA using PKCd(Phospho-Ser645) Antibody #11296.



Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using PKCd(Phospho-Ser645) Antibody #11296(left) or the same antibody preincubated with blocking peptide(right).

Background

This is calcium-independent, phospholipid-dependent, serine- and threonine-specific enzyme. PKC is activated by diacylglycerol which in turn phosphorylates a range of cellular proteins. PKC also serves as the receptor for phorbol esters, a class of tumor promoters. May play a role in antigen-dependent control of B-cell function. Phosphorylates MUC1 in the C-terminal and regulates the interaction between MUC1 and beta-catenin. Kei Sakamoto, et,al. (2003) Am J Physiol Endocrinol Metab ; 285: E1081 - E1088.

Ling Zhang, et,al. (2004) J. Biol. Chem ; 279: 28315 - 28319.

Kristof Van Kolen et,al. (2006) FEBS J ; 273: 1843 - 1854.

Martin Villalba, et,al. (2002) J. Cell Biol ; 157: 253.

Published Papers

M. Uenoyama, S. Ogata, K. Nakanishi el at., Protein kinase C mRNA and protein expressions in hypobaric hypoxia-induced cardiac hypertrophy in rats., Acta Physiol., 198:431B°C440(2010) PMID:19995357

Note: This product is for in vitro research use only and is not intended for use in humans or animals.