CREB(Phospho-Ser142) Antibody

Catalog No: #11300

Package Size: #11300-1 50ul #11300-2 100ul #11300-4 25ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Description


Product Name	CREB(Phospho-Ser142) Antibody
Host Species	Rabbit
Clonality	Polyclonal
Purification	Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates.
	Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho
	specific antibodies were removed by chromatogramphy using non-phosphopeptide.
Applications	WB IHC
Species Reactivity	Hu Ms Rt
Specificity	The antibody detects endogenous level of CREB only when phosphorylated at serine142.
Immunogen Type	Peptide-KLH
Immunogen Description	Peptide sequence around phosphorylation site of serine 142(D-L-S(p)-S-D) derived from Human CREB.
Target Name	CREB
Modification	Phospho-Ser142
Modification Other Names	Phospho-Ser142 CREB
	·
Other Names	CREB
Other Names Accession No.	CREB Swiss-Prot: P16220; NCBI Gene ID: 1385; NCBI mRNA: NM_004379.3; NCBI Protein: NP_004370.1
Other Names Accession No. SDS-PAGE MW	CREB Swiss-Prot: P16220; NCBI Gene ID: 1385; NCBI mRNA: NM_004379.3; NCBI Protein: NP_004370.1 43KD
Other Names Accession No. SDS-PAGE MW Concentration	CREB Swiss-Prot: P16220; NCBI Gene ID: 1385; NCBI mRNA: NM_004379.3; NCBI Protein: NP_004370.1 43KD 1.0mg/ml
Other Names Accession No. SDS-PAGE MW Concentration	CREB Swiss-Prot: P16220; NCBI Gene ID: 1385; NCBI mRNA: NM_004379.3; NCBI Protein: NP_004370.1 43KD 1.0mg/ml Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02%

Application Details


Predicted MW: 43kd

Immunohistochemistry: 1:50~1:100

Images



Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using CREB (Phospho-Ser142) Antibody (#11300).



Western blot analysis of extracts from 293 cells, treated with calf intestinal phosphatase (CIP), using CREB(Phospho-Ser142) Antibody #11300.

Background

Phosphorylation-dependent transcription factor that stimulates transcription upon binding to the DNA cAMP response element (CRE), a sequence present in many viral and cellular promoters. Transcription activation is enhanced by the TORC coactivators which act independently of Ser-133 phosphorylation. Involved in different cellular processes including the synchronization of circadian rhythmicity and the differentiation of adipose cells.

Published Papers

Tomasz Boczek, Anna Kozaczuk, Bozena Ferenc el at., Gene expression pattern in PC12 cells with reduced PMCA2 or PMCA3 isoform: selective up-regulation of calmodulin and neuromodulin, Mol Cell Biochem, 360(1-2):89-102(2012)

PMID:21912933

Note: This product is for in vitro research use only and is not intended for use in humans or animals.