# Cdc25B(Phospho-Ser149) Antibody

Catalog No: #11553

Package Size: #11553-1 50ul #11553-2 100ul #11553-4 25ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

_		4.5
	escri	ntion
$\boldsymbol{\nu}$	COUL	puon


Product Name	Cdc25B(Phospho-Ser149) Antibody	
Host Species	Rabbit	
Clonality	Polyclonal	
Purification	Antibodies were produced by immunizing rabbits with synthetic peptide and KLH conjugates. Antibodies were	
	purified by affinity-chromatography using epitope-specific peptide.	
Applications	WB	
Species Reactivity	Ms	
Specificity	The antibody detects endogenous level of Cdc25B protein only when phosphorylated at serine 149.	
Immunogen Type	Peptide-KLH	
Immunogen Description	Peptide sequence around phosphorylation site of Serine 149 (F-R-S(p)-L-P) derived from Mouse Cdc25B.	
Target Name	Cdc25B	
Modification	Phospho-Ser149	
Other Names	CDC25B	
Accession No.	Swiss-Prot: Q9DBN8NCBI Protein: NP_001104545.1	
Concentration	1.0mg/ml	
Formulation	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02%	
	sodium azide and 50% glycerol.	
Storage	Store at -20°C for long term preservation (recommended). Store at 4°C for short term use.	

### **Application Details**


Predicted MW: 62kd

Western blotting: 1:500~1:1000

### **Images**



Western blot analysis of extracts from 3T3 cells untreated(lane 1) or treated with Hydroxyurea(lane 2) using Cdc25B(Phospho-Ser149) Antibody #11553.



Western blot analysis of extracts from HT29 cells, treated with PMA or calf intestinal phosphatase (CIP), using Cdc25B (Phospho-Ser149) Antibody #11553.

## Background

Cdc25B is a member of the CDC25 family of phosphatases. CDC25B activates the cyclin dependent kinase CDC2 by removing two phosphate groups and it is required for entry into mitosis. CDC25B shuttles between the nucleus and the cytoplasm due to nuclear localization and nuclear export signals. The protein is nuclear in the M and G1 phases of the cell cycle and moves to the cytoplasm during S and G2. CDC25B has oncogenic properties, although its role in tumor formation has not been determined. Multiple transcript variants for this gene exist.

Li G, et al. (2010) Retrovirology.7:59.

Wang Z,et al. (2010) BMC Cancer.10:233.

Lucci MA,et al. (2010) Cell Oncol.32(5-6):361-72.

#### **Published Papers**

Jianying Xiao, Chao Liu, Junjie Hou el at., Ser149 Is Another Potential PKA Phosphorylation Target of Cdc25B in G2/M Transition of Fertilized Mouse Eggs., THE JOURNAL OF BIOLOGICAL CHEMISTRY, 286(12):10356B°C10366(2011)

PMID:21212267

Note: This product is for in vitro research use only and is not intended for use in humans or animals.