

## STAT2(Ab-690) Antibody

Catalog No: #21536



Package Size: #21536-1 50ul #21536-2 100ul #21536-4 25ul

Orders: [order@signalwayantibody.com](mailto:order@signalwayantibody.com)Support: [tech@signalwayantibody.com](mailto:tech@signalwayantibody.com)

## Description

Product Name	STAT2(Ab-690) Antibody
Host Species	Rabbit
Clonality	Polyclonal
Purification	Antibodies were produced by immunizing rabbits with synthetic peptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific peptide.
Applications	WB IHC
Species Reactivity	Hu
Specificity	The antibody detects endogenous level of total STAT2 protein.
Immunogen Type	Peptide-KLH
Immunogen Description	Peptide sequence around aa.688~692 (R-K-Y-L-K) derived from Human STAT2.
Target Name	STAT2
Other Names	P113; ISGF-3; STAT113
Accession No.	Swiss-Prot: P52630NCBI Protein: NP_005410.1
Concentration	1.0mg/ml
Formulation	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg <sup>2+</sup> and Ca <sup>2+</sup> ), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Storage	Store at -20°C for long term preservation (recommended). Store at 4°C for short term use.


## Application Details

Predicted MW: 113kd


Western blotting: 1:500~1:1000

Immunohistochemistry: 1:50~1:100

## Images



Western blot analysis of extracts from HeLa cells using STAT2(Ab-690) Antibody #21536 and the same antibody preincubated with blocking peptide.



Immunohistochemical analysis of paraffin-embedded human lung carcinoma tissue using STAT2(Ab-690) Antibody #21536(left) or the same antibody preincubated with blocking peptide(right).

## Background

STAT2 encoded by this gene is a member of the STAT protein family. In response to cytokines and growth factors, STAT family members are phosphorylated by the receptor associated kinases, and then form homo- or heterodimers that translocate to the cell nucleus where they act as transcription activators. In response to interferon (IFN), this protein forms a complex with STAT1 and IFN regulatory factor family protein p48 (ISGF3G), in which this protein acts as a transactivator, but lacks the ability to bind DNA directly. Transcription adaptor P300/CBP (EP300/CREBBP) has been shown to interact specifically with this protein, which is thought to be involved in the process of blocking IFN- $\alpha$  response by adenovirus.

Fu, X. et al. (1992) *Biochemistry* 89, 7840-7843.

Park, C. et al. (2000) *Immunity* 13, 795-804.

Horvath, C. et al. (1996) *Molecular and Cellular Biology* 16, 6957-6964.

Imprata, T. et al. (1994) *Proc. Natl. Acad. Sci. USA* 91, 4776-4780.

Note: This product is for in vitro research use only and is not intended for use in humans or animals.