## FLIP Antibody

Catalog No: #24045



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com


$\overline{}$			
	escr	TO	tion
$\boldsymbol{L}$	COUL	ıv	เเบเ

Product Name	FLIP Antibody	
Host Species	Rabbit	
Clonality	Polyclonal	
Purification	Affinity chromatography purified via peptide column	
Applications	E WB ICC	
Species Reactivity	Hu	
Specificity	Antibody recognizes human and murine FLIPgamma and FLIPdelta but not FLIPalpha and FLIPbeta.	
Immunogen Type	Peptide	
Immunogen Description	Raised against a peptide corresponding to amino acids 191 to 209 at C-terminus of human FLIP form2.	
Target Name	FLIP	
Other Names	FLIPs, FLIPg, d	
Accession No.	AAC51623	
Formulation	Supplied in PBS containing 0.02% sodium azide.	
Storage	Can be stored at -20°C, stable for one year. As with all antibodies care should be taken to avoid repeated	
	freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.	


## **Application Details**

Predicted MW: 35 kd and 25 kd

## **Images**



Western blot analysis of FLIP in total cell lysates from HeLa (H), Jurkat (J), THP-1 (T), A431 (A), K562 (K) and NIH3T3 (N) cells with FLIP antibody at 1:1000 dilution.



Immunocytochemistry staining of HeLa cells using FLIP antibody at 2 ug/mL.

## Background

Apoptosis is related to many diseases and induced by a family of cell death receptors and their ligands. Cell death signals are transduced by death domain (DD)-containing adapter molecules and members of the ICE/CED-3 protease family. Caspases-8 (FLICE) and -10 (FLICE2) are two pivotal members in the ICE/CED-3 protease family. FLICE-inhibitory proteins were identified in virus and human and designated v-FLIPs and c-FLIPs, respectively. The human FLIPs were also cloned by several labs independently and termed Casper, I-FLICE, FLAME-1, CASH, CLARP and usurpin. FLIP contains two death effector domains (DEDs) and a caspase-like domain. FLIP interacts with adapter protein FADD and caspase-8 and 10, and potently inhibits apoptosis induced by all known death receptors CD95, DR3, TRAIL-R and TNFR1. Four splice variants of c-FLIPs have been identified and termed FLIPalpha, beta, gamma, and delta, respectively.

Note: This product is for in vitro research use only and is not intended for use in humans or animals.