AIF Antibody

Catalog No: #24095



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com


## Description

Product Name	AIF Antibody
Host Species	Rabbit
Clonality	Polyclonal
Purification	Ion exchange chromatography purified
Applications	E WB ICC
Species Reactivity	Ни
Immunogen Type	Peptide
Immunogen Description	Raised against a peptide corresponding to amino acids near the amino terminus of mature human AIF.
Target Name	AIF
Accession No.	O95831
Formulation	Supplied in PBS containing 0.02% sodium azide.
Storage	Can be stored at -20°C, stable for one year. As with all antibodies care should be taken to avoid repeated
	freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.


## **Application Details**

Predicted MW: 67 kd

## Images



Western blot analysis of AIF in K562 cell lysate with AIF antibody at 1 ug/mL.



Immunocytochemistry of AIF in Jurkat cells with AIF antibody at 2 ug/mL.

## Background

Apoptosis is characterized by several morphological nuclear changes including chromatin condensation and nuclear fragmentation. These changes are triggered by the activation of members of caspase family, caspase activated DNase, and several novel proteins. A novel gene, the product of which causes chromatin condensation and DNA fragmentation, was recently identified, cloned, and designated apoptosis inducing factor (AIF). Like the critical molecules, cytochrome c and caspase-9, in apoptosis, AIF localizes in mitochondria. AIF translocates to the nucleus when apoptosis is induced and induces mitochondria to release the apoptogenic proteins cytochrome c and caspase-9. AIF induces chromatin condensation and DNA fragmentation, which are the hallmarks of apoptosis, of the isolated nucleus and the nucleus in live cells by microinjection. AIF is highly conserved between human and mouse and widely expressed.

Note: This product is for in vitro research use only and is not intended for use in humans or animals.